# Construction of double Grothendieck polynomials of classical types using Id-Coxeter algebras

Anatol N. Kirillov\* and Hiroshi Naruse\*\*

2015.04.30

Research Institute for Mathematical Sciences<sup>\*</sup>, RIMS, Kyoto University, Sakyo-ku, 606-8502 Japan and The Kavli Institute for Physics and Mathematics of the Universe<sup>\*</sup>, IPMU, 5-1-5 Kashiwanoha, Kashiwa, 277-8583 Japan. kirillov@kurims.kyoto-u.ac.jp

Graduate School of Education, University of Yamanashi\*\*, 4-4-37 Takeda, Kofu, Yamanashi, 400-8510 Japan. hnaruse@yamanashi.ac.jp

### Abstract

We construct double Grothendieck polynomials of classical types which are equivalent to the polynomials defined in [15] and compare with [14].

## 1 Introduction

Let G be a semisimple Lie group,  $B \subset G$  be a Borel subgroup of  $G, T \subset B$  be a maximal torus in  $B, \mathcal{F} := G/B$  and  $W := N_G(T)/T$  be the corresponding flag variety and the Weyl group. Let  $\ell$  be the rank of G.

According to the famous Borel theorem, the cohomology ring  $H^*(G/B, \mathbb{Q})$  is isomorphic to the quotient  $\mathbb{Q}[x_1, \ldots, x_\ell]/J_\ell$ , where  $x_i := c_1(L_i) \in H^2(G/B, \mathbb{Q})$ ,  $i = 1, \ldots, \ell$ , and  $c_1(L_i)$  denotes the first Chern class of the standard line bundle  $L_i$  over the flag variety in question,  $J_\ell$  stands for the ideal generated by the fundamental invariants of positive degree associated with the Weyl group W.

To our best knowledge the first systematic and complete treatment of the Schubert Calculus has been done by I.N. Bernstein, I.M. Gelfand and S.I. Gelfand [2] and independently, by M. Demazure [5] in the beginning of 70's of the last century. A Schubert polynomial  $\overline{\mathfrak{S}_w}(X_\ell)$ ,  $\ell = rk(G)$ , corresponding to an element w of the Weyl group W, by definition is a polynomial which expresses the Poincaré dual class of the homology class of the Schubert variety  $X_w := \overline{BwB/B} \subset G/B$  in terms of the Borel generators  $x_i, 1 \leq i \leq \ell$ , in the cohomology ring of the flag variety  $\mathcal{F}$ . Therefore by the very definition, a Schubert polynomial  $\overline{\mathfrak{S}_w}(X)$  is defined only modulo the ideal  $J_\ell$ .

Hence it is an interesting problem: does there exist "natural representative" of a Shubert polynomial  $\overline{\mathfrak{S}_w}(X_\ell)$  in the ring  $\mathbb{Q}[x_1,\ldots,x_\ell]$  with "nice" combinatorial, algebraic and geometric properties ?

For the type  $A_{n-1}$  flag varieties A. Lascoux and M.-P. Schütsenberger constructed a family of double Schubert polynomials  $\mathfrak{S}_w(X_{n-1}, Y_{n-1})$ ,  $w \in S_n$  with several "nice" properties, when we set  $\mathfrak{S}_w(X_{n-1}) := \mathfrak{S}_w(X_{n-1}, 0)$ , such as

1.  $\mathfrak{S}_w(X_{n-1})$  is a representative of the Schubert class  $X_w$  corresponding to  $w \in S_n$ , that is  $\mathfrak{S}_w(X_{n-1}) \equiv \overline{\mathfrak{S}}_w(X_{n-1}) \pmod{J_{n-1}}$ ,

2. (Compatibility conditions)

$$\partial_i^{(x)} \mathfrak{S}_w(X_{n-1}, Y_{n-1}) = \begin{cases} \mathfrak{S}_{ws_i}(X_{n-1}, Y_{n-1}) & \text{if } l(ws_i) = \ell(w) - 1, \\ 0 & \text{otherwise} \end{cases}$$

$$\partial_i^{(y)} \mathfrak{S}_w(X_{n-1}, Y_{n-1}) = \begin{cases} \mathfrak{S}_{s_i w}(X_{n-1}, Y_{n-1}) & \text{if } l(s_i w) = \ell(w) - 1, \\ 0 & \text{otherwise} \end{cases}$$

- 3.  $\mathfrak{S}_w(X_{n-1}, Y_{n-1})$  has nonnegative integer coefficients,
- 4.  $\mathfrak{S}_w(X_{n-1}, Y_{n-1})$  is stable,
- 5.  $\mathfrak{S}_w(X_{n-1}, Y_{n-1})$  satisfies the vanishing conditions, that is

$$\mathfrak{S}_w(-v(Y_{n-1}), Y_{n-1}) = 0$$
, unless  $w \leq v$ 

with respect to the Bruhat order  $\leq$  on the symmetric group  $S_n$ ,

6. the structural constants for the multiplication of Schubert polynomials  $\mathfrak{S}_w(X_{n-1})$ ,  $w \in S_n$ , coincide with the triple intersections numbers of Schubert varieties.

A new approach to the theory of type A Schubert polynomials which is based on the study of the type A nil–Coxeter algebras, has been initiated by S. Fomin and R. Stanley. The basic idea of that approach is to consider and study the generating function of all Schubert polynomials simultaneously, namely, to treat the following generating function

$$\mathfrak{S}(X_{n-1}) = \sum_{w \in S_n} \mathfrak{S}_w(X_{n-1})u_w,$$

where  $u_w$  denotes the standard linear basis in the nil-Coxeter algebra  $NC_n$ .

An unexpected and deep result discovered in [9] is that in the algebra  $NC_n[x_1, \ldots, x_{n-1}]$ the polynomial  $\mathfrak{S}_n(X_{n-1})$  is completely factorizable in the product of linear factors. The basic tool to prove the factorizability property is the usage of the Yang–Baxter relation among the elements  $h_i(x) = 1 + xu_i$  in the algebra  $NC_n[x, y]$ , namely

$$(1+xu_i)(1+(x+y)u_{i+1})(1+yu_i) = (1+yu_{i+1})(1+(x+y)u_i)(1+xu_{i+1}).$$
 (1)

The main consequence of the Yang–Baxter relation (1) is that the polynomials  $A_k(x) = h_{n-1}(x)h_{n-2}(x)\dots h_k(x)$ , commute, namely

$$[A_k(x), A_k(y)] = 0.$$

Now one can prove, [9], [8] that

$$\mathfrak{S}(X_{n-1}) = \sum_{w \in S_n} \mathfrak{S}_w(X_{n-1})u_w = A_1(x_1)A_2(x_2)\dots A_{n-1}(x_{n-1}).$$

This approach can be applied to a construction of type A double Schubert polynomials, Grothendieck and double Grothendieck polynomials, which originally had been introduced by A. Lascoux and M.-P. Schützenberger.

Construction of "good" representatives for the Schubert polynomials corresponding to the flag varieties of classical types B, C, D was initiated by S. Billey and M.Haiman [3] and independently by S. Fomin and A. N. Kirillov, [7]. In [7] the authors extended an algebrocombinatorial approach to a definition and study of the type A Schubert and Grothendieck polynomials to the case of those of types B and C. But it also works for type D as well. The key tool in a construction of the aforementioned polynomials is a unitary exponential solution to the quantum Yang–Baxter equations ([22]) with values in the NiCoxeter algebras of types B, C, D correspondingly. The exponential solution to the quantum Yang–Baxter equation associated with nilCoxeter algebra  $NC(R), R := A_{n-1}, B_n, C_n, D_n$ , allows to construct a family of elements  $R_i(x) \in NC(R)[x], i = 1, \ldots, rk(R)$  such that

$$R_i(x)R_i(y) = R_i(y)R_i(x), i = 1, \dots, rk(R).$$

The elements  $R_i(x_1), \ldots, R_i(x_\ell), i = 1, \ldots, \ell := rk(R)$ , are building blocks in the construction of the generating function for all Schubert polynomials corresponding to the flag variety associated with the root system R.

Now in order to ensure the coherency conditions one needs to specify the action of simple transpositions of the corresponding Weyl group on the ring of polynomials  $\mathbb{Q}[x_1, \ldots, x_\ell]$ . In [7] and [15] the authors have chosen the *standard* action of the Weyl group on the cohomology ring of the corresponding flag variety G/B. Namely,

$$s_0(x_1) = -x_1, s_0(x_i) = x_i, \text{ if } i \ge 1, (types B, C),$$
  
 $s_1^2 = -s_1, s_1(x_i) = x_i \text{ if } i \ge 2, (type D).$ 

Based on these (= standard !) choice of the action of the simple transpositions, the divided difference operators (resp. *isobaric* ones) are defined uniquely. It is easy to see [7] that for root systems of types B, C, D it is impossible to find "good" representatives for the Schubert classes which satisfy the properties 2, 3, 6 listed above. Nevertheless in [7] the authors introduce the so called Schubert polynomials of the second kind with nice combinatorial properties including those 3, 4, 5, 6, and therefore suitable for computation of the triple intersection numbers for Schubert polynomials of classical type, the main Problem of the Schubert Calculus, see [7] for details.

As for a construction of certain representatives the double Schubert and  $\beta$ -Grothendieck polynomials of classical types B, C, D, the author of [15] has used the following observation: if the a family of polynomials  $S_w(X), w \in W$  has "wanted properties" with respect to variables X, the polynomials

$$S_w(X,Y) := \sum_{\substack{u,v \in W, uv = w \\ \ell(uv) = \ell(u) + \ell(v)}} S_{u^{-1}}(Y) S_v(X)$$

also will have "wanted properties" with respect to the set of variables X and Y. Based on this observation and using the Schubert and Grothendieck polynomials introduced in [6], the author of [15] has introduced a family of polynomials depending on two sets of variable X and Y having nice combinatorial properties including among others, that 3, 4, 5, 6 listed above. One example of such polynomials is the triple  $\beta$ -Grothendieck polynomial  $\mathfrak{G}_w^W(X,Y,Z), w \in W$ , where W stands for the Weyl group of classical type B, C, or D. Indeed, let W be of type B, C, D, one can start with the W-type Schubert/Grothendieck expression of the second kind  $S^W(Z,X) := \sqrt{H^W(Z)}\mathfrak{G}^A(X)$ , have been introduced for the Schubert polynomials in [3] for Schubert polynomials of types W, [7] for W = B, C types Schubert polynomials, [16] for Schubert/Grothendieck case. According to an observation mentioned above, a "good" candidate for the <u>double</u> Schubert/Grothendieck expression of type W is

$$S^{W}(Y,T,Z,X) := S^{W}(-Y,-T)^{-1}S^{W}(Z,X) = \mathfrak{G}^{A}(-Y)^{-1}\sqrt{H^{W}(T)H^{W}(Z)}\mathfrak{G}^{A}(X).$$

To deduce this equality we have used the following facts:

$$\left[H^{W}(T), H^{W}(Z)\right] = 0, H^{W}(-T)^{-1} = H^{W}(T).$$

Finally, one can restrict the generating function  $S^W(Y, T, Z, X)$  on the diagonal T = Zand come to the following expression for the generating function of a double Schubert/Grothendieck polynomials of type W

$$S^W(Y, Z, X) = \mathfrak{G}^A(-Y)^{-1}H^W(Z)\mathfrak{G}^A(X).$$

Another algero-geometric interpretation of the generating function  $S^W(Y, Z, X)$  has been obtained in [12].

Advantage of the algebro-combinatorial approach is, for example, a possibility to define, among others, a plactic versions of polynomials  $\mathfrak{G}_w^W(X,Y)$ ,  $\mathfrak{S}_w^W(X,Y)$  and their generalizations, see [16] for the case of root systems of type A

In [3] the authors used non-standard action of Weyl group on the ring of supersymmetric functions of infinite number of variables  $\Gamma = (\mathbb{Z}[x_1, x_2, \ldots])^{SS}$  and define another family of Schubert polynomials.

In [12] the second author *et al.* studied the double Schubert polynomials of type B, C, D using localization map of equivariant cohomology. For K-theory there is analogous map and the image has the so called Goresky-Kottwitz-MacPherson property [10]. As mentioned for the case of Grassmannians in [14], the Schubert classes can be characterized by recurrence relations.(c.f. §6.)

## 2 Definitions and Notations

In this paper W = W(X) is a Weyl group of type X = A, B, C, D.  $I^X$  is the set of simple reflections in W(X). We index the simple reflections by the same notation as in [12] §3.2. In particular, for type B and C,  $s_0$  corresponds to the left most node of the Dynkin diagram with the relation  $(s_0s_1)^4 = 1$  and  $(s_0s_i)^2 = 1$  for  $i \ge 2$ . For type  $D, s_1 := s_0s_1s_0$ and we consider W(D) as the subgroup of W(B) generated by  $s_1, s_1, \ldots$ 

Following [7], we prepare some notations. Let  $\beta$  be an indeterminate. We define operations  $\oplus$  and  $\ominus$  as follows.

$$x \oplus y := x + y + \beta xy, x \ominus y := (x - y)/(1 + \beta y).$$

We also use the convention that

$$\bar{x} := \ominus x = -\frac{x}{1+\beta x}.$$

Then we have  $x \oplus \overline{x} = 0$ . For a Weyl group W with the set S of Coxeter generators, we define Id-Coxeter algebra as follows.

### **Definition 1.** (*Id-Coxeter algebra*)

Id-Coxeter algebra  $Id_{\beta}(W)$  for W is a  $\mathbb{Z}[\beta]$  algebra with generators  $u_i$  for each  $s_i \in S$  and relations as follows.

$$u_i u_j u_i \cdots = \beta u_i,$$

$$\underbrace{u_i u_j u_i \cdots}_{m_{i,j} \text{ termes}} = \underbrace{u_j u_i u_j \cdots}_{m_{i,j} \text{ termes}} \quad \text{if } m_{i,j} \text{ is the order of } s_i s_j.$$

For each  $s_i \in I^X$ , we define divided-difference operator  $\pi_i^{(a)}$  and  $\psi_i^{(a)}$  with respect to the variables  $a = (a_1, a_2, ...)$  as follows. Assume that  $R \supset \mathbb{Z}[\beta]$  is a ring with a group action of W(X). We define the action of W(X) on  $R[a, \bar{a}] := R[a_1, a_2, ..., \bar{a}_1, \bar{a}_2, ...]$  as follows.

**Definition 2.** The action of  $s_i \in I^X$  on the variables  $a_1, a_2, \ldots, \bar{a}_1, \bar{a}_2, \ldots$ 

- If  $i \ge 1$ ,  $s_i(a_i) = a_{i+1}, s_i(a_{i+1}) = a_i, s_i(\bar{a}_i) = \bar{a}_{i+1}, s_i(\bar{a}_{i+1}) = \bar{a}_i$ , and  $s_i(a_k) = a_k, s_i(\bar{a}_k) = \bar{a}_k$  for  $k \ne i, i+1$ .
- $s_0(a_1) = \bar{a}_1$ ,  $s_0(\bar{a}_1) = a_1$ , and  $s_0(a_k) = a_k$ ,  $s_0(\bar{a}_k) = \bar{a}_k$  for k > 1.
- $s_{\hat{1}}(a_1) = \bar{a}_2, s_{\hat{1}}(a_2) = \bar{a}_1, s_{\hat{1}}(\bar{a}_1) = a_2, s_{\hat{1}}(\bar{a}_2) = a_1, \text{ and } s_{\hat{1}}(a_k) = a_k, s_{\hat{1}}(\bar{a}_k) = \bar{a}_k \text{ for } k > 2.$

We write the induced action on  $R[a, \bar{a}]$  by  $s_i^{(a)}$ . Divided difference operators  $\pi_i^{(a)}$  and  $\psi_i^{(a)}$  are defined as follows. For  $f \in R[a, \bar{a}] = R[a_1, a_2, ..., \bar{a}_1, \bar{a}_2, ...]$ ,

$$\pi_i^{(a)}(f) := \frac{f - (1 + \beta \alpha_i(a)) s_i^{(a)}(f)}{\alpha_i(a)} \text{ and } \psi_i^{(a)} := \pi_i^{(a)} + \beta,$$

where  $\alpha_i(a)$  is the element in  $\mathbb{Z}[\beta][a, \bar{a}]$  corresponding to the root  $\alpha_i$ , i.e.  $\alpha_i(a) = a_i \oplus \bar{a}_{i+1}$ for  $i = 1, 2, ..., \alpha_0^B(a) = \bar{a}_1, \alpha_0^C(a) = \bar{a}_1 \oplus \bar{a}_1$  and  $\alpha_1(a) = \bar{a}_1 \oplus \bar{a}_2$ .

(Formally we can think as  $\alpha_i(a) = \frac{e^{\beta \alpha_i} - 1}{\beta}$ . c.f. [6])

**Proposition 1.** We have the following relations of operators.  $\pi_i^2 = -\beta \pi_i, \ \psi_i^2 = \beta \psi_i \ \text{for all } s_i \in I^X.$   $\underbrace{\pi_i \pi_j \pi_i \cdots}_{m_{i,j} \text{ termes}} = \underbrace{\pi_j \pi_i \pi_j \cdots}_{m_{i,j} \text{ termes}}, \ \underbrace{\psi_i \psi_j \psi_i \cdots}_{m_{i,j} \text{ termes}} = \underbrace{\psi_j \psi_i \psi_j \cdots}_{m_{i,j} \text{ termes}} \ \text{if } m_{i,j} \ \text{is the order of } s_i s_j.$ 

We can check the relations by direct calculations.

The explicit form of 
$$\psi_i^{(a)}$$
 is as follows,  
 $\psi_i^{(a)}(F) = \frac{s_i^{(a)}F - F}{a_{i+1} \ominus a_i}$  for  $i \ge 1$ ,  
 $\psi_{0,B}^{(a)}(F) = \frac{s_0^{(a)}F - F}{a_1}$ ,  $\psi_{0,C}^{(a)}(F) = \frac{s_0^{(a)}F - F}{a_1 \oplus a_1}$  and  $\psi_1^{(a)}(F) = \frac{s_1^{(a)}F - F}{a_1 \oplus a_2}$ .

Similarly we can define divided difference operators  $\pi_i^{(b)}$  and  $\psi_i^{(b)}$  corresponding to the variables  $b_1, b_2, \dots$ 

#### 3 **Basic Properties**

Let  $h_i(x) := 1 + xu_i$ . Then it follows that  $h_i(x)h_i(y) = h_i(x \oplus y)$ .

Lemma 1. (Yang-Baxter relation)  

$$h_i(x)h_j(y) = h_j(y)h_i(x)$$
 $m_{i,j} = 2$   
 $h_i(x)h_j(x \oplus y)h_i(y) = h_j(y)h_i(x \oplus y)h_j(x)$ 
 $m_{i,j} = 3$ 

$$h_i(x)h_j(x\oplus y)h_i(x\oplus y\oplus y)h_j(y) = h_j(y)h_i(x\oplus y\oplus y)h_j(x\oplus y)h_i(x) \quad m_{i,j} = 4$$

These can be proved by direct calculations.

### Definition 3.

$$\begin{aligned} A_i^{(n)}(x) &:= h_{n-1}(x)h_{n-2}(x)\cdots h_i(x) \quad (i=1,2,...,n-1) \\ F_n^B(x) &:= A_1^{(n)}(x) h_0(x) A_1^{(n)}(\bar{x})^{-1} \\ &= h_{n-1}(x)h_{n-2}(x)\cdots h_1(x)h_0(x)h_1(x)\cdots h_{n-2}(x)h_{n-1}(x) \\ F_n^C(x) &:= A_1^{(n)}(x) h_0(x)^2 A_1^{(n)}(\bar{x})^{-1} \\ &= h_{n-1}(x)h_{n-2}(x)\cdots h_1(x)h_0(x)^2h_1(x)\cdots h_{n-2}(x)h_{n-1}(x) \\ F_n^D(x) &:= A_2^{(n)}(x) h_1(x) h_1(x) A_2^{(n)}(\bar{x})^{-1} \\ &= h_{n-1}(x)\cdots h_2(x)h_1(x)h_1(x)h_2(x)\cdots h_{n-1}(x) \end{aligned}$$

### Lemma 2.

mma 2.  
(1) 
$$A_i^{(n)}(x)A_i^{(n)}(y) = A_i^{(n)}(y)A_i^{(n)}(x)$$
  
(2)  $F_n^X(x)F_n^X(y) = F_n^X(y)F_n^X(x)$  for  $X = B, C, D$   
(3)  $F_n^X(x)F_n^X(\bar{x}) = 1$ 

Note that from (1) we have  $A_i^{(n)}(x)A_i^{(n)}(y)^{-1} = A_i^{(n)}(y)^{-1}A_i^{(n)}(x)$  and  $A_i^{(n)}(x)^{-1}A_i^{(n)}(y)^{-1} = A_i^{(n)}(y)^{-1}A_i^{(n)}(x)^{-1}$ .

Proof.

(1) For the case i = n - 1 is trivial. By reverse induction on i, we can assume i < n - 1 and  $A_{i+1}^{(n)}(x)A_{i+1}^{(n)}(y) = A_{i+1}^{(n)}(y)A_{i+1}^{(n)}(x)$ . Then  $A_i^{(n)}(x)A_i^{(n)}(y) = A_{i+1}^{(n)}(x)h_i(x)A_{i+1}^{(n)}(y)h_i(y)$ 

$$= A_{i+1}^{(n)}(x)A_{i+1}^{(n)}(y)h_{i+1}(\bar{y})h_{i}(x)h_{i+1}(y)h_{i}(y \ominus x)h_{i}(x)$$

$$= A_{i+1}^{(n)}(x)A_{i+1}^{(n)}(x)h_{i+1}(\bar{y})h_{i+1}(y \ominus x)h_{i}(y)h_{i+1}(x)h_{i}(x)$$

$$= A_{i+1}^{(n)}(y)A_{i+1}^{(n)}(x)h_{i+1}(\bar{x})h_{i}(y)h_{i+1}(x)h_{i}(x)$$

$$= A_{i+1}^{(n)}(y)A_{i+2}^{(n)}(x)h_{i}(y)h_{i+1}(x)h_{i}(x)$$

$$= A_{i+1}^{(n)}(y)A_{i+2}^{(n)}(x)h_{i+1}(x)h_{i}(x)$$

$$= A_{i+1}^{(n)}(y)A_{i+2}^{(n)}(x)h_{i+1}(x)h_{i}(x)$$

$$= A_{i+1}^{(n)}(y)A_{i+2}^{(n)}(x)h_{i+1}(x)h_{i}(x)$$

(2) Using Lemma 3.1 and (1) we can show the equalities as follows. For X = B,

$$\begin{split} F_n^B(x)F_n^B(y) &= A_1^{(n)}(x)h_0(x)A_1^{(n)}(\bar{x})^{-1}A_1^{(n)}(y)h_0(y)A_1^{(n)}(\bar{y})^{-1} \\ &= A_1^{(n)}(x)h_0(x)A_1^{(n)}(y)A_1^{(n)}(\bar{x})^{-1}h_0(y)A_1^{(n)}(\bar{y})^{-1} \\ &= A_1^{(n)}(x)A_1^{(n)}(y)h_1(\bar{y})h_0(x)h_1(y)h_1(x)h_0(y)h_1(\bar{x})A_1^{(n)}(\bar{x})^{-1}A_1^{(n)}(\bar{y})^{-1} \\ &= A_1^{(n)}(y)A_1^{(n)}(x)h_1(\bar{y})h_0(x)h_1(y)h_1(x)h_0(y)h_1(\bar{x})A_1^{(n)}(\bar{y})^{-1}A_1^{(n)}(\bar{x})^{-1} \\ &= A_1^{(n)}(y)A_2^{(n)}(x)h_1(x\oplus\bar{y})h_0(x)h_1(x\oplus y)h_0(y)h_1(\bar{x}\oplus y)A_2^{(n)}(\bar{y})^{-1}A_1^{(n)}(\bar{x})^{-1} \\ &= A_1^{(n)}(y)A_2^{(n)}(x)h_0(y)h_1(x\oplus y)h_0(x)A_2^{(n)}(\bar{y})^{-1}A_1^{(n)}(\bar{x})^{-1} \\ &= A_1^{(n)}(y)h_0(y)A_1^{(n)}(x)A_1^{(n)}(\bar{y})^{-1}h_0(x)A_1^{(n)}(\bar{x})^{-1} \\ &= A_1^{(n)}(y)h_0(y)A_1^{(n)}(\bar{y})^{-1}A_1^{(n)}(x)h_0(x)A_1^{(n)}(\bar{x})^{-1} \\ &= F_n^B(y)F_n^B(x) \end{split}$$

Similar arguments with appropriate modifications will give X = C, D cases. The essential equalities to be used are

 $h_1(x \oplus \bar{y})h_0(x \oplus x)h_1(x \oplus y)h_0(y \oplus y)h_1(\bar{x} \oplus y) = h_0(y \oplus y)h_1(x \oplus y)h_0(x \oplus x) \text{ and } \\ h_2(x \oplus \bar{y})h_1(x)h_{\hat{1}}(x)h_2(x \oplus y)h_1(y)h_{\hat{1}}(y)h_2(\bar{x} \oplus y) = h_1(y)h_{\hat{1}}(y)h_2(x \oplus y)h_1(x)h_{\hat{1}}(x).$ (3) This esentially follows by the relation  $h_i(x)h_i(\bar{x}) = 1.$ 

## 4 $\beta$ -super symmetric functions

**Definition 4.**  $\beta$ -super symmetric function is a symmetric function which satisfies the following property.

 $f(t, \bar{t}, x_3, ..., x_n) = f(0, 0, x_3, ..., x_n)$  for every t.

### Remark

The  $\beta$ -supersymmetric property is translated to usual supersymmetricity by the change of variables  $x_i$  to  $\frac{e^{\beta x_i} - 1}{\beta}$ .

Let  $SS_{\beta}(x_1, \ldots, x_n) := \{ f \in \mathbb{Z}[\beta][x_1, \ldots, x_n] \mid f : \beta$ -supersymmetric  $\}$  and set  $SS_{\beta}(x) := \lim_{x \to \infty} SS_{\beta}(x_1, \ldots, x_n).$ 

 $\int^{n} SS_{\beta}(x)$  is the ring of  $\beta$ -supersymmetric functions and we denote it as  $\Gamma'_{\beta}(x)$ . If  $\beta = 0$  this becomes the the ring of supersymmetric functions  $\Gamma'$ .

## 4.1 K-theoretic Schur functions $GP_{\lambda}(x), GQ_{\lambda}(x)$

In [14]  $\beta$ -supersymmetric functions  $GP_{\lambda}(x), GQ_{\lambda}(x)$  are defined. Let  $b_1, b_2, ...$  be indeterminates, and set  $[x|b]^k = (x \oplus b_1) \cdots (x \oplus b_k)$  and  $[[x|b]]^k = (x \oplus x)(x \oplus b_1) \cdots (x \oplus b_{k-1})$ .

Let  $SP_n$  be the set of strict partitions of length at most n. i.e.  $\lambda = (\lambda_1 > \lambda_2 > \cdots > \lambda_r > 0)$  such that  $r \leq n$ .

**Definition 5.** (Ikeda-Naruse [14]) For a strict partition  $\lambda \in SP_n$ ,

$$GP_{\lambda}(x_1, \dots, x_n | b) := \frac{1}{(n-r)!} \sum_{w \in S_n} w \left( \prod_{1 \le i \le r} \left( [x_i | b]^{\lambda_i} \prod_{i < j \le n} \frac{x_i \oplus x_j}{x_i \ominus x_j} \right) \right)$$
$$GQ_{\lambda}(x_1, \dots, x_n | b) := \frac{1}{(n-r)!} \sum_{w \in S_n} w \left( \prod_{1 \le i \le r} \left( [[x_i | b]]^{\lambda_i} \prod_{i < j \le n} \frac{x_i \oplus x_j}{x_i \ominus x_j} \right) \right)$$

where  $w \in S_n$  acts  $x_1, \ldots, x_n$  as permutation of indices.

We also define

$$GP_{\lambda}(x_1, \dots, x_n) := GP_{\lambda}(x_1, \dots, x_n|0), \ GQ_{\lambda}(x_1, \dots, x_n) := GQ_{\lambda}(x_1, \dots, x_n|0),$$
$$GP_{\lambda}(x) := \lim_{\leftarrow n} GP_{\lambda}(x_1, \dots, x_n) \text{ and } GQ_{\lambda}(x) := \lim_{\leftarrow n} GQ_{\lambda}(x_1, \dots, x_n|b).$$
$$GP_{\lambda}(x|b) := \lim_{\leftarrow n} GP_{\lambda}(x_1, \dots, x_{2n}|b) \text{ and } GQ_{\lambda}(x|b) := \lim_{\leftarrow n} GQ_{\lambda}(x_1, \dots, x_n|b).$$

Examples.

 $GP_1(x_1,\ldots,x_n) = x_1 \oplus x_2 \oplus \cdots \oplus x_n.$  $GQ_1(x_1,\ldots,x_n) = (x_1 \oplus x_1) \oplus (x_2 \oplus x_2) \oplus \cdots \oplus (x_n \oplus x_n).$ 

### Lemma 3.

- (1)  $GP_{\lambda}(x_1, \ldots, x_n)$  and  $GQ_{\lambda}(x_1, \ldots, x_n)$  are  $\beta$ -supersymmetric functions.
- (2)  $\{GP_{\lambda}(x_1,\ldots,x_n)\}_{\lambda\in SP_n}$  forms a basis of  $SS_{\beta}(x_1,\ldots,x_n)$  over  $\mathbb{Z}[\beta]$ .

(3) Let  $SS^C_{\beta}(x_1, \ldots, x_n)$  be the  $\mathbb{Z}[\beta]$ -subspace of  $SS_{\beta}(x_1, \ldots, x_n)$  spanned by  $GQ_{\lambda}(x_1, \ldots, x_n)(\lambda \in SP_n)$ . Then  $\{GQ_{\lambda}(x_1, \ldots, x_n)\}_{\lambda \in SP_n}$  forms a basis of  $SS^C_{\beta}(x_1, \ldots, x_n)$  over  $\mathbb{Z}[\beta]$ .

Proof.

(1) follows from the definition.

(2) and (3) follows from the fact for corresponding properties for usual Schur P, Q-functions.

**Remark 1.** We remark that the definition of  $\beta$ -supersymmetry and the polynomials  $GP_{\lambda}, GQ_{\lambda}$  can be generalized in more general setting such as algebraic cobordism [20]. We are planning to study the details elsewhere. (cf.[21])

### Lemma 4. ([14])

 $GP_{\lambda}(x|b)$  and  $GQ_{\lambda}(x|b)$  are characterized by (left) divided difference relations and initial conditions. i.e.

$$\pi_i^{(b)} GX_{\lambda}(x|b) = \begin{cases} GX_{\lambda^{(i)}}(x|b) & \text{if } s_i\lambda < \lambda \\ -\beta \ GX_{\lambda}(x|b) & \text{if } s_i\lambda \ge \lambda \end{cases}$$

and

### $GX_{\emptyset}(x|b) = 1$

where  $GB_{\lambda}(x|b) = GP_{\lambda}(x|0,b), GC_{\lambda}(x|b) = GQ_{\lambda}(x|b), GD_{\lambda}(x|b) = GP_{\lambda}(x|b).$ 

See [14] Theorem 6.1 and Theorem 7.1.

# 4.2 Stable symmetric functions $\mathcal{F}_w^X(x_1,...,x_n)$

**Definition 6.** For X = B, C, D, we define

$$F_n^X(x_1, x_2, \dots, x_n) := \prod_{i=1}^n F_n^X(x_i) \text{ and } F_\infty^X(x) := \lim_{\leftarrow n} F_n^X(x_1, x_2, \dots, x_n).$$

We also define  $\mathcal{F}_w^X(x_1, ..., x_n)$  and  $\mathcal{F}_w^X(x)$  by the following expression.

$$F_n^X(x_1, x_2, \dots, x_n) = \sum_{w \in W_n^X} \mathcal{F}_w^X(x_1, \dots, x_n) u_w , F_\infty^X(x) = \sum_{w \in W^X} \mathcal{F}_w^X(x) u_w$$

**Lemma 5.** For each  $w \in W_n^X$ ,  $\mathcal{F}_w^X(x_1, x_2, \ldots, x_n)$  is a  $\beta$ -supersymmetric function.

Proof.

This follows from Lemma 3.3(2) and (3).

**Lemma 6.** (0) For  $X = B, C, D, \mathcal{F}_{w^{-1}}^X(x_1, x_2, \ldots, x_n) = \mathcal{F}_w^X(x_1, x_2, \ldots, x_n)$ . (1) For X = B or  $D, \mathcal{F}_w^X(x_1, x_2, \ldots, x_n)$  can be expanded in  $GP_\lambda(x_1, x_2, \ldots, x_n)$  with coefficients in  $\mathbb{Z}[\beta]$ .

(2) For a (maximal) Grassmannian element  $w \in W_n^X$ ,

$$\mathcal{F}_{w}^{B}(x_{1}, x_{2}, \dots, x_{n}) = GP_{\lambda_{B}(w)}(x_{1}, x_{2}, \dots, x_{n})$$
$$\mathcal{F}_{w}^{C}(x_{1}, x_{2}, \dots, x_{n}) = GQ_{\lambda_{C}(w)}(x_{1}, x_{2}, \dots, x_{n})$$
$$\mathcal{F}_{w}^{D}(x_{1}, x_{2}, \dots, x_{n}) = GP_{\lambda_{D}(w)}(x_{1}, x_{2}, \dots, x_{n})$$

where  $\lambda_B(w), \lambda_C(w), \lambda_D(w)$  are strict partitions corresponding to w. cf. [14]

Proof.

(0) This follows from the symmetry of  $F_n^X$ .

- (1) This follows from Lemma 4.3 (2).
- (2) This follows from Proposition 6.

**Remark 2.** We state conjecture that the coefficients in the expansion of (1) is positive. This will be a consequence of K-theory analogue of "transition equation" for type B, C, D. (cf. [12])

Example  $\mathcal{F}_{s_0}^B(x_1, \dots, x_n) = GP_1(x_1, \dots, x_n)$   $\mathcal{F}_{s_0}^C(x_1, \dots, x_n) = GQ_1(x_1, \dots, x_n)$  $\mathcal{F}_{s_1}^D(x_1, \dots, x_n) = GP_1(x_1, \dots, x_n)$ 

**Proposition 2.** (Compatible sequence formula) cf. ([3],[7]) For  $w \in W_n^X$ , we have

$$\mathcal{F}_{w}^{B}(x_{1},...,x_{n}) = \sum_{\tilde{a}\in\tilde{R}(w)} \sum_{\tilde{b}\in C^{B}(\tilde{a})} \beta^{\ell(\tilde{a})-\ell(w)} 2^{|\tilde{b}|-\gamma(\tilde{a},\tilde{b})-o^{B}(\tilde{a})} x_{\tilde{b}}$$
$$\mathcal{F}_{w}^{C}(x_{1},...,x_{n}) = \sum_{\tilde{a}\in\tilde{R}(w)} \sum_{\tilde{b}\in C^{C}(\tilde{a})} \beta^{\ell(\tilde{a})-\ell(w)} 2^{|\tilde{b}|-\gamma(\tilde{a},\tilde{b})} x_{\tilde{b}}$$
$$\mathcal{F}_{w}^{D}(x_{1},...,x_{n}) = \sum_{\tilde{a}\in\tilde{R}(w)} \sum_{\tilde{b}\in C^{D}(\tilde{a})} \beta^{\ell(\tilde{a})-\ell(w)} 2^{|\tilde{b}|-\gamma(\tilde{a},\tilde{b})-o^{D}(\tilde{a})} x_{\tilde{b}},$$

where we used the following notations.

R(w) is the set of sequence of indices  $\tilde{a} = (\tilde{a}_1, \tilde{a}_2, \dots, \tilde{a}_\ell)$  such that  $u_{s_{\tilde{a}_1}} \cdots u_{s_{\tilde{a}_\ell}} = u_w$ .  $\ell(\tilde{a})$  is the length  $\ell$  of the sequence  $\tilde{a}$ .

 $C^{B}(\tilde{a}) = C^{C}(\tilde{a})$  is the set of compatible sequences  $\tilde{b}$  with respect to  $\tilde{a}$ , i.e.  $\tilde{b} = (1 \leq \tilde{b}_{1} \leq \tilde{b}_{1} \leq \cdots \leq \tilde{b}_{\ell(\tilde{a})} \leq n)$  such that  $\tilde{a}_{i-1} \leq \tilde{a}_{i} \geq \tilde{a}_{i+1} \implies \tilde{b}_{i-1} < \tilde{b}_{i+1}$ .

 $C^{D}(\tilde{a})$  is the set of compatible sequence for the flattened word  $\tilde{\tilde{a}}$  of  $\tilde{a}$  with further properties that if  $\tilde{a}_{i} = \tilde{a}_{i+1} = 1$  or  $\tilde{a}_{i} = \tilde{a}_{i+1} = \hat{1}$  then  $\tilde{b}_{i} < \tilde{b}_{i+1}$ . Note that the flattened word  $\tilde{\tilde{a}}$  is obtained from  $\tilde{a}$  by replacing  $\hat{1}$  with 1. cf.[3].

 $o^B(\tilde{a})$  is the number of appearance of 0's in  $\tilde{a}$ .  $o^D(\tilde{a})$  is the total number of appearance of 1 and  $\hat{1}$  in  $\tilde{a}$ .  $|\tilde{b}|$  is the number of distinct  $\tilde{b}_i$ 's.  $\gamma(\tilde{a}, \tilde{b}) := \#|\{i|\tilde{a}_i = \tilde{a}_{i+1} \text{ and } \tilde{b}_i = \tilde{b}_{i+1}\}|.$   $x_{\tilde{b}} := x_{\tilde{b}_1} x_{\tilde{b}_2} \cdots x_{\tilde{b}_\ell}$  for  $\tilde{b} = (\tilde{b}_1 \dots, \tilde{b}_\ell).$ Proof. This follows essentially from the expansion of the definit

This follows essentially from the expansion of the defining generating function. Example.

type D, n = 2 case  $w = [\overline{1}, \overline{2}, 3] = s_1 s_1$  $\tilde{b} = (1, 1)$  is a compatible sequence for  $\tilde{a} = (1, 1), (1, 1).$  $\tilde{b} = (1, 2)$  is a compatible sequence for  $\tilde{a} = (1, 1), (1, 1).$  
$$\begin{split} \tilde{b} &= (2,2) \text{ is a compatible sequence for } \tilde{a} &= (1,\hat{1}), (\hat{1},1). \\ \tilde{b} &= (1,1,2) \text{ is a compatible sequence for } \tilde{a} &= (1,\hat{1},1), (\hat{1},1,\hat{1}), (1,\hat{1},\hat{1}), (\hat{1},1,1). \\ \tilde{b} &= (1,2,2) \text{ is a compatible sequence for } \tilde{a} &= (1,\hat{1},1), (\hat{1},1,\hat{1}), (1,1,\hat{1}), (\hat{1},\hat{1},1). \\ \tilde{b} &= (1,1,2,2) \text{ is a compatible sequence for } \tilde{a} &= (1,\hat{1},1), (\hat{1},1,\hat{1}), (\hat{1},1,\hat{1}), (\hat{1},1,\hat{1},1). \\ \tilde{a} &= (1,\hat{1},1,\hat{1}), (\hat{1},1,\hat{1},1), (\hat{1},1,1,\hat{1}), (1,\hat{1},\hat{1},1). \end{split}$$

There are no other compatible sequences and the sum of the terms becomes  $\mathcal{F}^{D}_{s_{1}s_{1}}(x_{1}, x_{2}) = x_{1}^{2} + 2x_{1}x_{2} + x_{2}^{2} + 2\beta x_{1}^{2}x_{2} + 2\beta x_{1}x_{2}^{2} + \beta^{2}x_{1}^{2}x_{2}^{2} = (x_{1} \oplus x_{2})^{2}.$ 

## 5 Main results

First we recall the type A Grothendieck polynomials [7].

We set  $G_{A_{n-1}}(a_1, ..., a_{n-1}) := A_1^{(n)}(a_1)A_2^{(n)}(a_2)\cdots A_{n-1}^{(n)}(a_{n-1}).$ Then for  $w \in S_n$ , we define  $\mathcal{G}_w^{A_{n-1}}(a)$  as the coefficient of  $u_w$ .

$$G_{A_{n-1}}(a_1, ..., a_{n-1}) = \sum_{w \in S_n} \mathcal{G}_w^{A_{n-1}}(a) u_w.$$

Furthermore, we can consider  $G_A(a) := \lim_{\leftarrow n} G_{A_{n-1}}(a_1, ..., a_{n-1})$  and get strongly stable polynomials  $\mathcal{G}_w^A(a)$  by

$$G_A(a) = \sum_{w \in S_\infty} \mathcal{G}_w^A(a) u_w.$$

Strongly stable means that if  $w \in S_n$  then  $\mathcal{G}_w^A(a) = \mathcal{G}_w^{A_{n-1}}(a)$  (which does not depend on n).

### 5.1 The first definition

**Definition 7.** We define for X = B, C or D,

$$G_n^X(a,b;x) := G_{A_{n-1}}(\bar{b}_1,...,\bar{b}_{n-1})^{-1}F_n^X(x)G_{A_{n-1}}(a_1,...,a_{n-1})$$

and define  $\mathcal{G}_{n,w}^X(a,b;x)$  as the coefficient of  $u_w$ .

$$G_n^X(a,b;x) = \sum_{w \in W_n^X} \mathcal{G}_{n,w}^X(a,b;x) u_w.$$

In this case  $\mathcal{G}_{n,w}^X(a,b;x) \in SS_\beta(x_1,\ldots,x_n)[a_1,\ldots,a_{n-1},b_1,\ldots,b_{n-1}].$ Furthermore, we can define  $\mathcal{G}_w^X(a,b;x)$  by

$$G_A(\bar{b})^{-1}F_\infty(x)G_A(a) = \sum_{w \in W^X} \mathcal{G}_w^X(a,b;x)u_w.$$

Then  $\mathcal{G}_w^X(a,b;x)$  has strong stability (cf. Proposition 5), and when we set  $\beta = 0$  this is the double Schubert polynomial defined in [12]. It is clear that if  $w \in W_n^X$  then  $\mathcal{G}_{n,w}^X(a,b;x) = \mathcal{G}_w^X(a,b;x_1,\ldots,x_n,0,0,\ldots)$ .

We will write  $w \cdot v = z$  (called Demazure product) if  $u_w u_v = \beta^{\ell(w) + \ell(v) - \ell(z)} u_z$ .

**Proposition 3.** For X = B, C, D and  $w \in W^X$ , we have

$$\mathcal{G}_w^X(a,b;x) = \sum_{(v_1,u,v_2)\in R(w)} \mathcal{G}_{v_1^{-1}}^A(b) \mathcal{F}_u^X(x) \mathcal{G}_{v_2}^A(a)$$

where  $R(w) = \{(v_1, u, v_2) \in S_{\infty} \times W^X \times S_{\infty} \mid v_1 \cdot u \cdot v_2 = w\}.$ 

**Definition 8.** The action of Weyl group  $W_n^X$  on  $SS_{\beta}^{(n)}(x) \otimes \mathbb{Z}[\beta][a, \bar{a}] \otimes \mathbb{Z}[\beta][b, \bar{b}]$  is derived from the action as follows. For  $f(x) \in SS_{\beta}(x)$ ,

$$s_0^{(a)} f(x) = f(a_1, x), \ s_0^{(b)} f(x) = f(b_1, x),$$
  
$$s_{\hat{1}}^{(a)} f(x) = f(a_1, a_2, x), \ s_{\hat{1}}^{(b)} f(x) = f(b_1, b_2, x).$$

These actions can be clarified by the change of variables explained in the second definition below (cf.§5.2 Remark 3).

### **Proposition 4.** We have

$$\pi_i^{(a)}G_n^X(a,b;x) = G_n^X(a,b;x)(u_i - \beta) \text{ and } \pi_i^{(b)}G_n^X(a,b;x) = (u_i - \beta)G_n^X(a,b;x)$$

N.B. These mean that

$$\pi_i^{(a)} \mathcal{G}_w^X(a,b;x) = \begin{cases} \mathcal{G}_{ws_i}^X(a,b;x) & \text{if } l(ws_i) = \ell(w) - 1, \\ -\beta \mathcal{G}_w^X(a,b;x) & \text{otherwise} \end{cases}$$

and

$$\pi_i^{(b)} \mathcal{G}_w^X(a,b;x) = \begin{cases} \mathcal{G}_{s_iw}^X(a,b;x) & \text{if } l(s_iw) = \ell(w) - 1, \\ -\beta \mathcal{G}_w^X(a,b;x) & \text{otherwise} \end{cases}$$

Proof.

We will prove  $\psi_i^{(a)} G_n^X(a,b;x) = G_n^X(a,b;x)u_i$ . Recall the explicit formula of  $\psi_i$  after the Prop. 2.2.

 $G_{A_{n-1}}(\bar{b})^{-1} \text{ is invariant for the action of } s_i^{(a)}, \ i \in I^X. \text{ For } i > 0, \ \psi_i^{(a)} F_n^X(x) = F_n^X(x)$ and  $\psi_i^{(a)} G_{A_{n-1}}(a) = G_{A_{n-1}}(a) u_i(\text{cf. [6]}), \text{ therefore } \psi_i^{(a)} F_n^X(x) G_{A_{n-1}}(a) = F_n^X(x) G_{A_{n-1}}(a) u_i(a)$ 

$$\psi_{0,B}^{(a)}(F_n^B(x)G_{A_{n-1}}(a)) = \frac{F_n^B(x)F^B(a_1)G_{A_{n-1}}(\bar{a}_1,a_2,\dots,a_{n-1}) - F_n^B(x)G_{A_{n-1}}(a)}{a_1} = F_n^B(x)G_{A_{n-1}}(a)u_0$$
  

$$\psi_{0,C}^{(a)}(F_n^C(x)G_{A_{n-1}}(a)) = \frac{F_n^C(x)H^C(a_1)G_{A_{n-1}}(\bar{a}_1,a_2,\dots,a_{n-1}) - F_n^C(x)G_{A_{n-1}}(a)}{a_1\oplus a_1} = F_n^C(x)G_{A_{n-1}}(a)u_0$$
  

$$\psi_{\hat{1}}^{(a)}(F_n^D(x)G_{A_{n-1}}(a)) = \frac{F_n^D(x)F^D(a_1,a_2)G_{A_{n-1}}(\bar{a}_2,\bar{a}_1,\dots,a_{n-1}) - F_n^D(x)G_{A_{n-1}}(a)}{a_1\oplus a_2} = F_n^D(x)G_{A_{n-1}}(a)u_{\hat{1}}$$
  
Similar arguments hold for the action of  $\psi_{\hat{1}}^{(b)}$ .

ents hold for the action of  $\psi_i$ 

**Proposition 5.** (strong stability)  $\hat{\mathcal{G}}_w^X(a,b;x)$  has strong stability i.e. if  $i_n: W_n^X \to W_{n+1}^X$  is the natural inclusion, then

$$\mathcal{G}_{i_n(w)}^{\scriptscriptstyle A}(a,b;x) = \mathcal{G}_w^{\scriptscriptstyle A}(a,b;x).$$

**Proposition 6.** (Grassmannian elements) For a Grassmannian element  $w \in W^X$ , we have the following equality.

$$\mathcal{G}_{w}^{B}(a,b;x) = GP_{\lambda_{B}(w)}(x|0,b)$$
$$\mathcal{G}_{w}^{C}(a,b;x) = GQ_{\lambda_{C}(w)}(x|b)$$
$$\mathcal{G}_{w}^{D}(a,b;x) = GP_{\lambda_{D}(w)}(x|b)$$

where  $\lambda_X(w)$  is the strict partition corresponding to  $w \in W^X$  (cf. [14]).

#### 5.2The second definition

As [7], we can use "change of variables" for  $x_i, i = 1, 2, \ldots$ 

$$F(x_i) = \sqrt{F(\bar{a}_i)F(\bar{b}_i)}$$

to define the double Grothendieck polynomial  $\mathcal{G}_w^{X_n}(a, b)$  with two sets of variables a, b.

**Remark 3.** As  $s_0^{(a)}(\sqrt{F(\bar{a}_1, \bar{a}_2, \ldots)}) = \sqrt{F(a_1, \bar{a}_2, \ldots)}$  and by the supersymmetric property of F, this is  $= \sqrt{F(a_1, a_1, \bar{a}_1, \bar{a}_2, \ldots)} = F(a_1)\sqrt{F(\bar{a}_1, \bar{a}_2, \ldots)}$ . This explains the action  $s_0^{(a)}(F(x)) = F(a_1, x)$  and  $s_0^{(b)}(F(x)) = F(b_1, x)$ . The action of  $s_1^{(a)}$  and  $s_1^{(b)}$  as well.

**Definition 9.** Let X = B, C, D. For  $w \in W_n^X$ , we define  $G_n^X(a)$  and  $G_n^X(a, b)$  as follows.

$$G_n^X(a) := \sqrt{F_n^X(\bar{a}_1, \dots, \bar{a}_n)} G_{A_{n-1}}(a) \quad and \quad G_n^X(a, b) := G_n^X(\bar{b})^{-1} G_n^X(a).$$

By expanding these in terms of  $u_w$ , we can define  $\mathcal{G}_{n,w}^X(a)$  and  $\mathcal{G}_{n,w}^X(a,b)$  by

$$G_n^X(a) = \sum_{w \in W_n^X} \mathcal{G}_{n,w}^X(a) u_w \quad and \quad G_n^X(a,b) = \sum_{w \in W_n^X} \mathcal{G}_{n,w}^X(a,b) u_w$$

**Remark 4.** This double Grothendieck polynomial  $\mathcal{G}_{n,w}^X(a,b)$  is essentially the same as defined in [15]. This has weak stability. i.e.  $\mathcal{G}_{n,w}^X = \mathcal{G}_{n+1,w}^X|_{a_{n+1}=b_{n+1}=0}$  for  $w \in W_n^X$ . But it doesn't have strong stability.

Note that for  $w \in W_n^X$ , then

$$\mathcal{G}_{n,w}^{X}(a) \in \mathbb{Q}[\beta][[a_{1},...,a_{n},\bar{a}_{1},...,\bar{a}_{n}]] \text{ and} \mathcal{G}_{n,w}^{X}(a,b) \in \mathbb{Q}[\beta][[a_{1},...,a_{n},\bar{a}_{1},...,\bar{a}_{n},b_{1},...,b_{n},\bar{b}_{1},...,\bar{b}_{n}]].$$

### Examples

$$\mathcal{G}^{B}_{2,s_{0}}(a,b) = \frac{\sqrt{1 + (\bar{a}_{1} \oplus \bar{a}_{2} \oplus \bar{b}_{1} \oplus \bar{b}_{2})\beta - 1}}{\beta} = \frac{\bar{a}_{1} \oplus \bar{a}_{2} \oplus \bar{b}_{1} \oplus \bar{b}_{2}}{2} - \beta \frac{(\bar{a}_{1} \oplus \bar{a}_{2} \oplus \bar{b}_{1} \oplus \bar{b}_{2})^{2}}{8} + \cdots \\
\mathcal{G}^{C}_{2,s_{0}}(a,b) = \bar{a}_{1} \oplus \bar{a}_{2} \oplus \bar{b}_{1} \oplus \bar{b}_{2}, \quad \mathcal{G}^{C}_{3,s_{0}}(a,b) = \bar{a}_{1} \oplus \bar{a}_{2} \oplus \bar{a}_{3} \oplus \bar{b}_{1} \oplus \bar{b}_{2} \oplus \bar{b}_{3}. \\
\mathcal{G}^{D}_{3,s_{1}}(a,b) = \frac{\sqrt{1 + (\bar{a}_{1} \oplus \bar{a}_{2} \oplus \bar{a}_{3} \oplus \bar{b}_{1} \oplus \bar{b}_{2} \oplus \bar{b}_{3})\beta - 1}{\beta}$$

**Proposition 7.** The following holds for X = B, C, D and  $i \in I_{X_n}$ .

$$\pi_i^{(a)} G_n^X(a, b) = G_n^X(a, b)(u_i - \beta)$$
$$\pi_i^{(b)} G_n^X(a, b) = (u_i - \beta) G_n^X(a, b)$$

Proof.

These are Prop. 5.3 with change of variables.

## 6 Identification with Schubert class

Let  $\mathcal{R}_{\beta}^{(b)} := \mathbb{Z}[\beta][b_1, \bar{b}_1, b_2, \bar{b}_2, \ldots]$ . *K*-theory Schubert classes are determined by the localization (Prop. 2.10 in [17]). And they are determined uniquely by either "right hand" recurrence ((2.12) in [17]) or "left hand" recurrence (Remark 2.3 in [17]).

(right recurrence)  $\mathcal{G}_e = 1$ , and  $\pi_i^{(a)} \mathcal{G}_w = \mathcal{G}_{ws_i}$  if  $ws_i < w$  and  $\pi_i^{(a)} \mathcal{G}_w = -\beta \mathcal{G}_w$  if  $ws_i > w$ .

(left recurrence)  $\mathcal{G}_e = 1$ , and  $\pi_i^{(b)} \mathcal{G}_w = \mathcal{G}_{s_iw}$  if  $s_i w < w$  and  $\pi_i^{(b)} \mathcal{G}_w = -\beta \mathcal{G}_w$  if  $s_i w > w$ . Therefore we can identify the polynomials  $\mathcal{G}_w^X(a, b; x)$  defined above as Schubert classes. In particular we have

**Theorem 1.** Assume 
$$\mathcal{G}_{u}^{X}(a,b;x)\mathcal{G}_{v}^{X}(a,b;x) = \sum_{w \in W^{X}} c_{u,v}^{w,X}(\beta)\mathcal{G}_{w}^{X}(a,b;x), \ c_{u,v}^{w,X}(\beta) \in \mathcal{R}_{\beta}^{(b)}.$$

Then  $c_{u,v}^{w,X}(\beta)|_{\beta=-1}$  is the generalized Littlewood-Richardson coefficient for equivariant Ktheory of type X. (b<sub>i</sub> is considered as  $1 - e^{t_i}$ .) **Remark 5.**  $c_{u,v}^w(0)$  is the generalized Littlewood-Richardson coefficient for equivariant cohomology if we replace  $b_i$  to  $-t_i$ . (cf. [12].)

Example  $\mathcal{G}^C_{s_0}(a,b;x)\mathcal{G}^C_{s_0}(a,b;x) = \mathcal{G}^C_{s_1s_0}(a,b;x) + \beta \mathcal{G}^C_{s_0s_1s_0}(a,b;x)$ 

## 7 Adjoint polynomials

The Grothendieck polynomial represents the K-theory Schubert class of the structure sheaf  $\mathcal{O}_{X_w}$  of the Schubert variety  $X^w = \overline{B_- w B/B} \subset X = G/B$ . We can also define the adjoint polynomials  $\mathcal{H}_{n,w}^X$ , for each  $w \in W_n^X$ , corresponding to the ideal sheaf  $\mathcal{O}_{X^w}(-\partial X^w)$ of boundary  $\partial X^w$  in  $X^w$ . cf. [11, 18]. The pairing  $\langle \cdot, \cdot \rangle : K_T(X) \otimes_{R(T)} K_T(X) \to R(T)$ is given by

$$\langle v_1, v_2 \rangle = \chi(X, v_1 \otimes v_2)$$
 where  $\chi(X, \mathcal{F}) = \sum_{p \ge 0} (-1)^p \mathrm{ch} \ H^p(X, \mathcal{F}).$ 

We define the relative adjoint polynomial  $\mathcal{H}_{w,v}^X$  for  $w \leq v$  by  $\mathcal{H}_{w,v}^X := \psi_{w^{-1}v}^{(a)}(\mathcal{G}_v^X)$ . The adjoint polynomial for  $w \in W_n^X$  is  $\mathcal{H}_{n,w}^X := \mathcal{H}_{w,w_0}^X$ , where  $w_0^{(n)}$  is the longest element in  $W_n^X$  (cf. [18]). These polynomials are no more stable but have similar properties as Grothendieck polynomials.

**Proposition 8.** For  $w \in W_n^X$ 

$$\begin{aligned} \mathcal{H}_{n,e}^{B} &= \prod_{1 \le i \le n-1} (1+\beta a_{i})^{n-i} \prod_{1 \le i \le n-1} (1+\beta b_{i})^{n-i} \prod_{1 \le i \le n} (1+\beta x_{i})^{2n-1} \\ \mathcal{H}_{n,e}^{C} &= \prod_{1 \le i \le n-1} (1+\beta a_{i})^{n-i} \prod_{1 \le i \le n-1} (1+\beta b_{i})^{n-i} \prod_{1 \le i \le n} (1+\beta x_{i})^{2n} \\ \mathcal{H}_{n,e}^{D} &= \prod_{1 \le i \le n-1} (1+\beta a_{i})^{n-i} \prod_{1 \le i \le n-1} (1+\beta b_{i})^{n-i} \prod_{1 \le i \le n} (1+\beta x_{i})^{2n-2} \end{aligned}$$

and

$$\mathcal{H}_{n,w}^X = (-1)^{\ell(w)} \mathcal{H}_e^{X_n} \overline{\mathcal{G}_{n,w}^X}$$

where  $\overline{\mathcal{G}_{n,w}^X} = \mathcal{G}_{n,w}^X(\overline{a}, \overline{b}; \overline{x}).$ 

We can derive these formula using generating functions. Let us define  $H_n^X(a,b;x)$  as

$$H_n^X(a,b;x) := \sum_{w \in W_n^X} (-1)^{\ell(w)} \mathcal{H}_{n,w}^X(a,b;x) u_w.$$

Then we get the following formula.

### Proposition 9.

$$H_n^X(a,b;x) = \mathcal{H}_{n,e}^X G_n^X(\bar{a},\bar{b};\bar{x}).$$

Actually we can show the following property.

**Proposition 10.** For  $s_i \in I_n^X$  we have

$$\pi_i^{(a)} H_n^X(a, b; x) = H_n^X(a, b; x)(-u_i)$$
  
$$\pi_i^{(b)} H_n^X(a, b; x) = (-u_i) H_n^X(a, b; x).$$

**Proposition 11.** (Interpolation formula) For  $F \in SS_{\beta} \otimes_{\mathbb{Z}[\beta]} \mathcal{R}_{\beta}^{(a)} \otimes_{\mathbb{Z}[\beta]} \mathcal{R}_{\beta}^{(b)}$ ,

$$F = \sum_{w \in W^X} (\psi_w^{(a)}(F)|_e) \ \mathcal{G}_w^X(a,b;x)$$

where the summation is infinite in general and  $|_e$  means the localization at e, i.e. take substitutions  $a_i = \bar{b}_i$  and  $x_i = 0$  for all i.

Corollary 1. The equivariant Littlewood-Richardson coefficient can be written as

$$c_{u,v}^{w,X}(\beta) = \psi_w^{(a)}(\mathcal{G}_u^X(a,b;x)\mathcal{G}_v^X(a,b;x))|_e.$$

Theorem 2.

$$\mathcal{G}_w^X(a,b;x) = \sum_{uv=w,u \le w} \mathcal{H}_{u,w}^X(\bar{c},b;0) \mathcal{G}_v^X(a,c;x)$$

There is also similar formula using second version of type B, C, D double Grothendieck polynomials.

## 8 Pipe dream formula

For type A Schubert/Grothendieck polynomials, it is well known that there is an explicit formula using pipe dream ([1, 6]). We can extend this to type B, C, D cases as follows. For this we recall the type A case formula. We give two descriptions, one in terms of excited Young diagrams [14], another one in terms of compatible sequences [16].

The type  $A_{n-1}$  double Grothendieck polynomials  $\mathcal{G}_w^{A_{n-1}}(a,b)$  are defined as follows.

$$G_{A_{n-1}}(\bar{b}_1, ..., \bar{b}_{n-1})^{-1}G_{A_{n-1}}(a_1, ..., a_{n-1}) = \sum_{w \in S_n} \mathcal{G}_w^{A_{n-1}}(a, b)u_w.$$

By using Yang-Baxter relations, the left hand side can be expressed as below

$$\prod_{j=n-1}^{1} \prod_{i=1}^{n-j} h_{i+j-1}(a_i \oplus b_j).$$
(2)

Therefore the expansion in terms of  $u_w$  produces the pipe dream formula.

1. Excited Young diagrams (EYD for short). Let N := n(n-1)/2 and

$$\Delta_n := (s_{n-1})(s_{n-2}, s_{n-1}) \cdots (s_1, s_2, \cdots, s_{n-1}) = (d_1, d_2, \dots, d_N)$$

Each term in the expansion of (2) corresponds to a subsequence of  $\Delta_n$ . Give  $w \in S_n$  with  $\ell(w) = \ell$ , let  $Rsub(\Delta_n, w)$  be the set of subsequences of  $\Delta_n$  each element of which gives a reduced expression of w. i.e.

$$Rsub(\Delta_n, w) := \{ (d_{j_1}, d_{j_2}, \dots, d_{j_\ell}) \mid 1 \le j_1 < j_2 < \dots < j_\ell \le N, d_{j_1} d_{j_2} \cdots d_{j_\ell} = w \}.$$

We will call  $D \in Rsub(\Delta_n, w)$  an extended EYD. For an extended EYD  $D = (d_{j_1}, d_{j_2}, \ldots, d_{j_\ell}) \in Rsub(\Delta_n, w)$ , we define the set B(D) of backward movable positions by, considering  $j_{\ell+1} := N + 1$ ,

$$B(D) := \{ d_j \mid j \le N, \exists p \text{ such that } j_p < j < j_{p+1}, d_{j_1} d_{j_2} \cdots d_{j_p} = (d_{j_1} d_{j_2} \cdots d_{j_p}) \cdot d_j \}.$$

We also define weight  $wt(d_k) = a_i \oplus b_j$  if k = (n-j)(n-j-1)/2 + i. Then we have the following extended EYD formula.

$$\mathcal{G}_{w}^{A_{n-1}}(a,b) = \sum_{D \in RSub(\Delta_{n},w)} Wt(D),$$

where

$$Wt(D) = \prod_{\Box \in D} wt(\Box) \times \prod_{\bigcirc \in B(D)} (1 + \beta wt(\bigcirc))$$

In the pipe dream diagram two patterns appear. One is  $\square$  which corresponds to the selected box  $\square$  in EYD configuration. The other case we put  $\square$  in the box. Each selected box  $\square$  in D corresponds to a word of the reduced expression of w inside  $\Delta_n$ . B(D) is the set of backward movable positions (cf. [14]).

Example: type  $A_3$ ,  $w = [3, 1, 4, 2] = s_2 s_3 s_1$ .



Example 2 
$$w = s_2 s_3 s_2 = s_3 s_2 s_3$$
  
One can show that  $\mathcal{G}_w^{A_{n-1}}(a,b)|_{a=1,b=0} = 5 + 5\beta + \beta^2$ .





 $\begin{array}{lll} Wt(D_1) &=& (a_2 \oplus b_2)(a_2 \oplus b_1)(a_3 \oplus b_1) \\ Wt(D_2) &=& (a_1 \oplus b_3)(a_2 \oplus b_1)(a_3 \oplus b_1)(1 + \beta(a_2 \oplus b_2)) \\ Wt(D_3) &=& (a_1 \oplus b_3)(a_1 \oplus b_2)(a_3 \oplus b_1)(1 + \beta(a_2 \oplus b_1)) \\ Wt(D_4) &=& (a_1 \oplus b_3)(a_1 \oplus b_2)(a_2 \oplus b_2)(1 + \beta(a_2 \oplus b_1))(1 + \beta(a_3 \oplus b_1)) \\ Wt(D_5) &=& (a_1 \oplus b_2)(a_2 \oplus b_2)(a_2 \oplus b_1)(1 + \beta(a_3 \oplus b_1)) \end{array}$ 

From these data we get  $\mathcal{G}_{s_2s_3s_2}^{A_3}(a,b) = Wt(D_1) + Wt(D_2) + Wt(D_3) + Wt(D_4) + Wt(D_5).$ 

Actually there is an algorithm to create all the extended EYD diagrams for a given  $w \in S_n$ . The algorithm is essentially written in [1]. Combinatorics related to extended EYD diagrams (including type B, C, D case) will be discussed elsewhere.

2. Compatible sequence.

Let T be a semistandard tableau and w(T) be the column reading word corresponding to the tableau T. Denote by R(T) (resp. IR(T))the set of words which are plactic (resp. idplactic) equivalent to w(T). Let  $\mathbf{a} = (a_1, \dots, a_n) \in R(T)$ , where n := |T| (resp.  $\mathbf{a} = (a_1, \dots, a_m) \in IR(T)$ , where  $m \ge |T|$ ).

**Definition 10.** ([8],[16]) (Compatible sequences  $\begin{cases} a \\ b \end{cases}$ )

Given a word  $\mathbf{a} \in R(T)$  (esp.  $\mathbf{a} \in IR(T)$ ), denote by  $C(\mathbf{a})$  (resp.  $IC(\mathbf{a})$ ) the set of sequences of positive integers, called compatible sequences,  $\mathbf{b} := (b_1 \leq b_2 \leq \cdots \leq b_m)$  such that

 $b_i \leq a_i, \text{ and if } a_i \leq a_{i+1}, \text{ then } b_i < b_{i+1}.$  (3)

Finally, define the set C(T) (resp. IC(T)) to be the union  $\bigcup C(\mathbf{a})$  (resp. the union  $\bigcup IC(\mathbf{a})$ ), where **a** runs over all words which are plactic (resp. idplactic) equivalent to the word w(T).

Example Take  $T = \frac{2}{3}$ <sup>3</sup>. The corresponding tableau word is w(T) = 323. We have  $R(T) = \{232, 323\}$  and  $IR(T) = R(T) \cup \{2323, 3223, 3232, 3233, 3323, 32323, \cdots\}$ . Moreover,

$$C(T) = \begin{cases} \mathbf{a} : 232 \ 323 \ 323 \ 323 \ 323 \ 323 \end{cases},$$
$$IC(T) = C(T) \bigcup \begin{cases} \mathbf{a} : 2323 \ 3223 \ 3223 \ 3223 \ 3223 \ 3223 \ 3232 \ 3233 \ 32323 \ 3233 \ 32323 \ 32323 \$$

From these data we get single Grothendieck polynomial

 $\mathcal{G}_{w}^{A_{n-1}}(a) = a_{1}a_{2}^{2} + a_{1}^{2}a_{2} + a_{1}^{2}a_{3} + a_{1}a_{2}a_{3} + a_{2}^{2}a_{3} + \beta(2a_{1}a_{2}^{2}a_{3} + 2a_{1}^{2}a_{2}a_{3} + a_{1}^{2}a_{2}^{2}) + \beta^{2}a_{1}^{2}a_{2}^{2}a_{3}.$ 

For type  $B_n$  or  $C_n$  case, we can rewrite the generating function of Definition 5.1 as follows.

$$\left(\prod_{j=n-1}^{1}\prod_{i=1}^{n-j}h_{i+j-1}(x_{n-i+1}\oplus b_j)\right)\left(\prod_{i=n}^{1}\prod_{j=n}^{i}h_{j-i}(x_{i,j}^X)\right)\left(\prod_{i=n-1}^{1}\prod_{j=1}^{n-i}h_{i+j-1}(x_i\oplus a_j)\right)$$
(4)

where  $x_{i,j}^X = x_i \oplus x_j$  if  $i \neq j$ ,  $x_{i,i}^B = x_i$  and  $x_{i,i}^C = x_i \oplus x_i$ . Comparing this to the type A case, we get the following formula.

**Proposition 12.** For  $w \in W(A_{n-1}) \subset W(B_n) = W(C_n)$ , we have  $\mathcal{G}^B_{n,w}(a,b;x) = \mathcal{G}^C_{n,w}(a,b;x) = \mathcal{G}^{A_{2n-1}}_{1^n \times w}(x_1, \dots, x_n, a_1, \dots, a_{n-1}, x_1, \dots, x_n, b_1, \dots, b_{n-1}).$ 

For type  $D_n$  case, we assume n = 2m an even integer, for odd n = 2m - 1 case we can get the formula by just erasing the last variable  $x_{2m} = 0$  for n = 2m case.

$$\left(\prod_{i=n}^{2}\prod_{j=i-1}^{1}h_{i+j-1}(x_{i}\oplus b_{j})\right)\left(\prod_{i=n-1}^{1}\prod_{j=n}^{i+1}h_{i,j}(x_{i}\oplus x_{j})\right)\left(\prod_{i=n-1}^{1}\prod_{j=1}^{n-i}h_{i+j-1}(x_{i}\oplus a_{j})\right)$$
(5)

where  $h_{i,j}(x_i \oplus x_j) := h_{j-i}(x_i \oplus x_j)$  if  $j - i \ge 2$ ,  $h_{i,i+1}(x_i \oplus x_{i+1}) := h_1(x_i \oplus x_{i+1})$  if i = odd, and  $h_{i,i+1}(x_i \oplus x_{i+1}) := h_1(x_i \oplus x_{i+1})$  if i = even.

If w is a maximal Grassmannian element of type  $B_n$ ,  $C_n$  or  $D_n$ , then the above gives the Excited Young diagram formula of [14] Th 9.2. Therefore the above give a generalization of the EYD formula.

Example

Type  $C_3$ ,  $w = [2, \bar{3}, 1] = s_2 s_1 s_2 s_0 s_1$ .



 $\times (1 + \beta(x_1 \oplus x_3))(1 + \beta(a_1 \oplus x_2))(1 + \beta(x_1 \oplus x_1))$ 

Example type  $D_4$ ,  $w = [\overline{2}, 4, \overline{1}, 3] = s_3 s_1 s_2$ 



### Conclusion

In the present paper we compare an algebro-combinatorial [15] and algebro- geometric [14] constructions of the double Schubert/Grothendieck polynomials of types B,C,D, and show that these two approaches give rise to essentially the same polynomial representatives for the Schubert/Grothendieck classes in the cohomology/K-theory rings of the types B,C and D full flag varieties correspondingly. The formulas obtained (4) and (5) lead to combinatorial descriptions of polynomials in questions in terms of either EYD, or compatible sequences, or set-valued tableaux [4].

We expect that after a certain change of Id-Coxeter algebra and replacing  $A \oplus B$  in our formulas (4) and (5) by F(A, B), where F(x, y) stands for the universal formal group law, we come to formal power series which have a suitable interpretations in the theory of algebraic cobordism [20] of flag varieties.

## References

- N.Bergeron and S.Billey, RC-Graphs and Schubert Polynomials, Experiment. Math. Vol. 2 (1993), 257–269.
- [2] I. N. Bernstein, I. M. Gelfand, and S. I. Gelfand, Schubert cells and cohomology of the spaces G/P, Russian Math. Surveys 28 (1973), no. 3, 1–26.
- [3] S. Billey and M. Haiman, Schubert polynomials for the classical groups, J. Amer. Math. Soc. 8 (1995), no. 2, 443–482.

- [4] A. Buch, A Littlewood–Richardson rule for the K-theory of Grassmannians, Acta. Math. 189 (2002), 2633–2640.
- [5] M. Demazure, Désingularisation des variétés de Schubert généralisées, Ann. Sci. Ecole Norm. Sup. 7 (1974), 53–88.
- [6] S.Fomin and A.N.Kirillov, Grothendieck polynomials and the Yang-Baxter equation. Proceedings of the Sixth Conference in Formal Power Series and Algebraic Combinatorics, DIMACS, 1994, pp. 183–190.
- [7] S.Fomin and A.N.Kirillov, Combinatorial  $B_n$ -analogues of Schubert polynomials. Trans. Amer. Math. Soc. 348 (1996), no. 9, 3591–3620.
- [8] S.Fomin and A.N. Kirillov, Yang-Baxter equation, symmetric functions and Grothendieck polynomials, preprint arXiv:hep-th/9306005.
- S.V. Fomin and R. Stanley, Schubert polynomials and the nil-Coxeter algebra, Adv. Math. 103 (1994), no. 2, 196–207.
- [10] M. Goresky, R Kottwitz, and R. MacPherson, Equivariant cohomology, Koszul duality, and the localization theorem, Invent. Math. 131 (1998) 25–83.
- [11] Graham and S. Kumar, On positivity in *T*-equivariant *K*-theory of flag varieties, Int. Math. Res. Not. IMRN 2008, Art. ID rnn 093.
- [12] T.Ikeda, L.Mihalcea and H.Naruse, Double Schubert polynomials for the classical groups, Adv. Math. 226 (2011), 840–886.
- [13] T.Ikeda and H.Naruse, Excited Young diagrams and equivariant Schubert calculus, Trans. Amer. Math. Soc. 361 (2009), no. 10, 5193–5221.
- [14] T.Ikeda and H.Naruse, K-theory analogue of factorial Schur P-,Q- functions, Adv. Math. 243 (2013), 22–66.
- [15] A.N.Kirillov, On Double Schubert and Grothendieck polynomials for classical groups, preprint (1999); update version arXiv:1504.0146.
- [16] A.N. Kirillov, Notes on Schubert, Grothendieck and Key polynomials, arXiv:1501.07337.
- [17] T.Lam, A.Schilling and M.Shimozono, K-Theory Schubert calculus of the affine Grassmannian, Compositio Math. 146 (2010), 811–852.

- [18] A.Lascoux, Anneau de Grothendieck de la variété de drapeaux, The Grothendieck Festschrift, Vol. III, 1–34, Progr. Math., 88, Birkhüser Boston, Boston, MA, 1990.
- [19] A. Lascoux and M.-P. Schützenberger, Polynômes de Schubert, C.R. Acad. Sci. Paris Sér. I Math., 294 (1982), 447–450.
- [20] M.Levin and F.Morel, Algebraic cobordism, Springer Monograph (2008).
- [21] M. Nakagawa and H. Naruse, Generalized (co)homology of the loop spaces of classical groups and the universal factorial Schur *P* and *Q*-functions, arXiv:1310.8008.
- [22] E.K. Sklyanin, L.A. Takhtadzhyan, L.D. Faddeev, Quantum inverse problem method.I, Theoretical and Mathematical Physics, Volume 40, Issue 2, pp.688–706,1979.