

Non-unitary holography

Daniel Grumiller

Institute for Theoretical Physics
TU Wien

IPMU, January 2015

H. Afshar, J. Aparicio, A. Bagchi, M. Bertin, B. Cvetkovic,
S. Detournay, S. Ertl, M. Gaberdiel, M. Gary, O. Hohm,
R. Jackiw, N. Johansson, E. Lopez, I. Papadimitriou,
R. Rashkov, I. Sachs, S. Stricker, D. Vassilevich, T. Zojer



Quoted from the workshop webpage:

1. How general is holography?

To what extent do (previous) lessons rely on the particular constructions used to date? Are they tied to stringy effects and to string theory in particular, or are they general lessons for quantum gravity?

Appetizer

Quoted from the workshop webpage:

1. How general is holography?

To what extent do (previous) lessons rely on the particular constructions used to date? Are they tied to stringy effects and to string theory in particular, or are they general lessons for quantum gravity?

Historically: holography intimately related to/derived from unitarity ('t Hooft '93, Susskind '94)

Appetizer

Quoted from the workshop webpage:

1. How general is holography?

To what extent do (previous) lessons rely on the particular constructions used to date? Are they tied to stringy effects and to string theory in particular, or are they general lessons for quantum gravity?

Historically: holography intimately related to/derived from unitarity ('t Hooft '93, Susskind '94)

Specific question addressed in this talk:

Does holography apply only to unitary theories?

Focus on very special but generic phenomenon

Leitmotif: critical points

Focus on very special but generic phenomenon

Leitmotif: critical points

Basic idea (generic):

$$(i\frac{d}{dt} - \omega_1)(i\frac{d}{dt} - \omega_2)\psi = 0$$

$\omega_{1,2}$ determined by parameters/coupling constants of theory

Two branches of solutions: $\psi_{1,2} \propto e^{-i\omega_{1,2} t}$

Focus on **very special** but **generic** phenomenon

Leitmotif: **critical points**

Basic idea (**generic**):

$$(i\frac{d}{dt} - \omega_1)(i\frac{d}{dt} - \omega_2)\psi = 0$$

$\omega_{1,2}$ determined by parameters/coupling constants of theory

Two branches of solutions: $\psi_{1,2} \propto e^{-i\omega_{1,2}t}$

Critical point through infinite fine tuning (**very special**): $\omega_2 \rightarrow \omega_1$

Focus on **very special** but **generic** phenomenon

Leitmotif: **critical points**

Basic idea (**generic**):

$$(i\frac{d}{dt} - \omega_1)(i\frac{d}{dt} - \omega_2)\psi = 0$$

$\omega_{1,2}$ determined by parameters/coupling constants of theory

Two branches of solutions: $\psi_{1,2} \propto e^{-i\omega_{1,2}t}$

Critical point through infinite fine tuning (**very special**): $\omega_2 \rightarrow \omega_1$

Branches degenerate and “**logarithmic**” branch emerges:

$$\psi^{\log} \propto \lim_{\omega_2 \rightarrow \omega_1} \frac{\psi_2 - \psi_1}{\omega_2 - \omega_1} \propto t e^{-i\omega_1 t} + \alpha \psi_1$$

Hamiltonian $H = i\frac{d}{dt}$ acquires **Jordan-block structure**

$$H \begin{pmatrix} \psi^{\log} \\ \psi_1 \end{pmatrix} = \begin{pmatrix} \omega_1 & 1 \\ 0 & \omega_1 \end{pmatrix} \begin{pmatrix} \psi^{\log} \\ \psi_1 \end{pmatrix}$$

Focus on **very special** but **generic** phenomenon

Leitmotif: **critical points**

Basic idea (**generic**):

$$(i\frac{d}{dt} - \omega_1)(i\frac{d}{dt} - \omega_2)\psi = 0$$

$\omega_{1,2}$ determined by parameters/coupling constants of theory

Two branches of solutions: $\psi_{1,2} \propto e^{-i\omega_{1,2}t}$

Critical point through infinite fine tuning (**very special**): $\omega_2 \rightarrow \omega_1$

Branches degenerate and “**logarithmic**” branch emerges:

$$\psi^{\log} \propto \lim_{\omega_2 \rightarrow \omega_1} \frac{\psi_2 - \psi_1}{\omega_2 - \omega_1} \propto te^{-i\omega_1 t} + \alpha \psi_1$$

Hamiltonian $H = i\frac{d}{dt}$ acquires **Jordan-block structure**

$$H \begin{pmatrix} \psi^{\log} \\ \psi_1 \end{pmatrix} = \begin{pmatrix} \omega_1 & 1 \\ 0 & \omega_1 \end{pmatrix} \begin{pmatrix} \psi^{\log} \\ \psi_1 \end{pmatrix}$$

Holographic version of **critical points**?

Outline

Gravity in three dimensions

Logarithmic CFTs

$\text{AdS}_3/\text{LCFT}_2$ correspondence

Generalizations

Outline

Gravity in three dimensions

Logarithmic CFTs

$\text{AdS}_3/\text{LCFT}_2$ correspondence

Generalizations

Motivations for studying gravity in 3 dimensions

► Quantum gravity

- Address conceptual issues of quantum gravity
- Black hole evaporation, information loss, black hole microstate counting, virtual black hole production, ...
- Integrable models: powerful tools in physics (Coulomb problem, Hydrogen atom, harmonic oscillator, ...)
- Models should be as simple as possible, but not simpler

► Gauge/gravity duality

- Deeper understanding of black hole holography
- $\text{AdS}_3/\text{CFT}_2$ correspondence best understood
- Quantum gravity via AdS/CFT? (Witten '07, Li, Song, Strominger '08)
- Applications to 2D condensed matter systems?
- Gauge gravity duality beyond standard AdS/CFT: warped AdS, asymptotic Schrödinger/Lifshitz, non-relativistic CFTs, flat space holography, higher spin holography, logarithmic CFTs, ...

► Physics

- Cosmic strings (Deser, Jackiw, 't Hooft '84, '92)
- Black hole analog systems in condensed matter physics (graphene, BEC, fluids, ...)

Topologically massive gravity

Deser, Jackiw & Templeton '82

$$I_{\text{TMG}} = \frac{1}{16\pi G} \int d^3x \sqrt{-g} \left[R + \frac{2}{\ell^2} + \frac{1}{2\mu} \varepsilon^{\lambda\mu\nu} \Gamma^\rho_{\lambda\sigma} (\partial_\mu \Gamma^\sigma_{\nu\rho} + \frac{2}{3} \Gamma^\sigma_{\mu\tau} \Gamma^\tau_{\nu\rho}) \right]$$

Properties:

- Parity odd gravitational Chern–Simons term with coupling μ

Topologically massive gravity

Deser, Jackiw & Templeton '82

$$I_{\text{TMG}} = \frac{1}{16\pi G} \int d^3x \sqrt{-g} \left[R + \frac{2}{\ell^2} + \frac{1}{2\mu} \varepsilon^{\lambda\mu\nu} \Gamma^\rho_{\lambda\sigma} (\partial_\mu \Gamma^\sigma_{\nu\rho} + \frac{2}{3} \Gamma^\sigma_{\mu\tau} \Gamma^\tau_{\nu\rho}) \right]$$

Properties:

- ▶ Parity odd gravitational Chern–Simons term with coupling μ
- ▶ Asymptotic AdS, warped AdS, Schrödinger/Lifshitz, $\text{AdS}_2 \times S^1$, flat, ?

Topologically massive gravity

Deser, Jackiw & Templeton '82

$$I_{\text{TMG}} = \frac{1}{16\pi G} \int d^3x \sqrt{-g} \left[R + \frac{2}{\ell^2} + \frac{1}{2\mu} \varepsilon^{\lambda\mu\nu} \Gamma^\rho_{\lambda\sigma} (\partial_\mu \Gamma^\sigma_{\nu\rho} + \frac{2}{3} \Gamma^\sigma_{\mu\tau} \Gamma^\tau_{\nu\rho}) \right]$$

Properties:

- ▶ Parity odd gravitational Chern–Simons term with coupling μ
- ▶ Asymptotic AdS, warped AdS, Schrödinger/Lifshitz, $\text{AdS}_2 \times S^1$, flat, ?
- ▶ Interesting scaling limits ($\mu \rightarrow \infty$, $\mu \rightarrow 0$, $\mu\ell \rightarrow 1$, $\ell \rightarrow \infty$, ...)

Topologically massive gravity

Deser, Jackiw & Templeton '82

$$I_{\text{TMG}} = \frac{1}{16\pi G} \int d^3x \sqrt{-g} \left[R + \frac{2}{\ell^2} + \frac{1}{2\mu} \varepsilon^{\lambda\mu\nu} \Gamma^\rho_{\lambda\sigma} (\partial_\mu \Gamma^\sigma_{\nu\rho} + \frac{2}{3} \Gamma^\sigma_{\mu\tau} \Gamma^\tau_{\nu\rho}) \right]$$

Properties:

- ▶ Parity odd gravitational Chern–Simons term with coupling μ
- ▶ Asymptotic AdS, warped AdS, Schrödinger/Lifshitz, $\text{AdS}_2 \times S^1$, flat, ?
- ▶ Interesting scaling limits ($\mu \rightarrow \infty$, $\mu \rightarrow 0$, $\mu\ell \rightarrow 1$, $\ell \rightarrow \infty$, ...)
- ▶ Black hole solutions, massive gravitons

Equations of motion:

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R - \frac{1}{\ell^2} g_{\mu\nu} + \frac{1}{\mu} C_{\mu\nu} = 0$$

Cotton tensor

$$C_{\mu\nu} = \frac{1}{2} \varepsilon_\mu^{\alpha\beta} \nabla_\alpha R_{\beta\nu} + (\mu \leftrightarrow \nu)$$

Outline

Gravity in three dimensions

Logarithmic CFTs

$\text{AdS}_3/\text{LCFT}_2$ correspondence

Generalizations

Conformal field theories in two dimensions

- ▶ Any CFT has conserved traceless energy momentum tensor (EMT)

$$T_{z\bar{z}} = 0 \quad T_{zz} = \mathcal{O}^L(z) \quad T_{\bar{z}\bar{z}} = \mathcal{O}^R(\bar{z})$$

Conformal field theories in two dimensions

- ▶ Any CFT has conserved traceless energy momentum tensor (EMT)

$$T_{z\bar{z}} = 0 \quad T_{zz} = \mathcal{O}^L(z) \quad T_{\bar{z}\bar{z}} = \mathcal{O}^R(\bar{z})$$

- ▶ 2- and 3-point correlators fixed by conformal Ward identities

$$\langle \mathcal{O}^L(z) \mathcal{O}^L(0) \rangle = \frac{c_L}{2z^4} \quad \langle \mathcal{O}^R(\bar{z}) \mathcal{O}^R(0) \rangle = \frac{c_R}{2\bar{z}^4}$$

Central charges $c_{L/R}$ determine key properties of CFT

Conformal field theories in two dimensions

- ▶ Any CFT has conserved traceless energy momentum tensor (EMT)

$$T_{z\bar{z}} = 0 \quad T_{zz} = \mathcal{O}^L(z) \quad T_{\bar{z}\bar{z}} = \mathcal{O}^R(\bar{z})$$

- ▶ 2- and 3-point correlators fixed by conformal Ward identities

$$\langle \mathcal{O}^L(z) \mathcal{O}^L(0) \rangle = \frac{c_L}{2z^4} \quad \langle \mathcal{O}^R(\bar{z}) \mathcal{O}^R(0) \rangle = \frac{c_R}{2\bar{z}^4}$$

Central charges $c_{L/R}$ determine key properties of CFT

- ▶ Minimal models:

$$c_{(p,q)} = 1 - 6 \frac{(p-q)^2}{pq} \quad \Rightarrow c_{(3,2)} = 0$$

Kac table only one entry: identity — only trivial $c = 0$ CFTs?

Conformal field theories in two dimensions

- ▶ Any CFT has conserved traceless energy momentum tensor (EMT)

$$T_{z\bar{z}} = 0 \quad T_{zz} = \mathcal{O}^L(z) \quad T_{\bar{z}\bar{z}} = \mathcal{O}^R(\bar{z})$$

- ▶ 2- and 3-point correlators fixed by conformal Ward identities

$$\langle \mathcal{O}^L(z) \mathcal{O}^L(0) \rangle = \frac{c_L}{2z^4} \quad \langle \mathcal{O}^R(\bar{z}) \mathcal{O}^R(0) \rangle = \frac{c_R}{2\bar{z}^4}$$

Central charges $c_{L/R}$ determine key properties of CFT

- ▶ Minimal models:

$$c_{(p,q)} = 1 - 6 \frac{(p-q)^2}{pq} \quad \Rightarrow \quad c_{(3,2)} = 0$$

Kac table only one entry: identity — only trivial $c = 0$ CFTs?

- ▶ Concrete $c = 0$ model exist where partition function is not trivial: percolation, self-avoiding polymers, $O(n)$ model in $n \rightarrow 0$ limit, systems with quenched disorder, ...

Conformal field theories in two dimensions

- Any CFT has conserved traceless energy momentum tensor (EMT)

$$T_{z\bar{z}} = 0 \quad T_{zz} = \mathcal{O}^L(z) \quad T_{\bar{z}\bar{z}} = \mathcal{O}^R(\bar{z})$$

- 2- and 3-point correlators fixed by conformal Ward identities

$$\langle \mathcal{O}^L(z) \mathcal{O}^L(0) \rangle = \frac{c_L}{2z^4} \quad \langle \mathcal{O}^R(\bar{z}) \mathcal{O}^R(0) \rangle = \frac{c_R}{2\bar{z}^4}$$

Central charges $c_{L/R}$ determine key properties of CFT

- Minimal models:

$$c_{(p,q)} = 1 - 6 \frac{(p-q)^2}{pq} \quad \Rightarrow \quad c_{(3,2)} = 0$$

Kac table only one entry: identity — only trivial $c = 0$ CFTs?

- Concrete $c = 0$ model exist where partition function is not trivial: percolation, self-avoiding polymers, $O(n)$ model in $n \rightarrow 0$ limit, systems with quenched disorder, ...
- Cannot be described by minimal (unitary) models

The $c = 0$ catastrophe

Primary field \mathcal{O}^M with conformal weights (h, \bar{h}) :

$$\langle \mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \rangle = \frac{A}{2z^{2h} \bar{z}^{2\bar{h}}}$$

OPE:

$$\mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \sim \frac{A}{2z^{2h} \bar{z}^{2\bar{h}}} \left(1 + \frac{2h}{c_L} z^2 \mathcal{O}^L(0) + \dots \right)$$

Problem: divergence for $c_L \rightarrow 0$

The $c = 0$ catastrophe

Primary field \mathcal{O}^M with conformal weights (h, \bar{h}) :

$$\langle \mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \rangle = \frac{A}{2z^{2h} \bar{z}^{2\bar{h}}}$$

OPE:

$$\mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \sim \frac{A}{2z^{2h} \bar{z}^{2\bar{h}}} \left(1 + \frac{2h}{c_L} z^2 \mathcal{O}^L(0) + \dots \right)$$

Problem: divergence for $c_L \rightarrow 0$

Possible resolutions in limit $c_L \rightarrow 0$:

- ▶ weight vanishes $h \rightarrow 0$
- ▶ normalization vanishes $A \rightarrow 0$
- ▶ other operator(s) arise with $h \rightarrow 2$, which cancel divergence

Focus on last possibility

Correlators in logarithmic conformal field theories (LCFTs)

Aghamohammadi, Khorrami & Rahimi Tabar '97; Kogan & Nichols '01; Rasmussen '04

Suppose now that primary has conformal weights $(2 + \varepsilon, \varepsilon)$:

$$\langle \mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \rangle = \frac{A}{2z^{4+2\varepsilon} \bar{z}^{2\varepsilon}}$$

Correlators in logarithmic conformal field theories (LCFTs)

Aghamohammadi, Khorrami & Rahimi Tabar '97; Kogan & Nichols '01; Rasmussen '04

Suppose now that primary has conformal weights $(2 + \varepsilon, \varepsilon)$:

$$\langle \mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \rangle = \frac{A}{2z^{4+2\varepsilon} \bar{z}^{2\varepsilon}}$$

Suppose that limits exist:

$$b_L := - \lim_{c_L \rightarrow 0} \frac{c_L}{\varepsilon} \neq 0 \quad A = -c_L + \mathcal{O}(c_L^2)$$

Correlators in logarithmic conformal field theories (LCFTs)

Aghamohammadi, Khorrami & Rahimi Tabar '97; Kogan & Nichols '01; Rasmussen '04

Suppose now that primary has conformal weights $(2 + \varepsilon, \varepsilon)$:

$$\langle \mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \rangle = \frac{A}{2z^{4+2\varepsilon} \bar{z}^{2\varepsilon}}$$

Suppose that limits exist:

$$b_L := - \lim_{c_L \rightarrow 0} \frac{c_L}{\varepsilon} \neq 0 \quad A = -c_L + \mathcal{O}(c_L^2)$$

Define \log operator:

$$\mathcal{O}^{\log} = \lim_{\varepsilon \rightarrow 0} \frac{\mathcal{O}^M - \mathcal{O}^L}{\varepsilon}$$

Correlators in logarithmic conformal field theories (LCFTs)

Aghamohammadi, Khorrami & Rahimi Tabar '97; Kogan & Nichols '01; Rasmussen '04

Suppose now that primary has conformal weights $(2 + \varepsilon, \varepsilon)$:

$$\langle \mathcal{O}^M(z, \bar{z}) \mathcal{O}^M(0, 0) \rangle = \frac{A}{2z^{4+2\varepsilon} \bar{z}^{2\varepsilon}}$$

Suppose that limits exist:

$$b_L := - \lim_{c_L \rightarrow 0} \frac{c_L}{\varepsilon} \neq 0 \quad A = -c_L + \mathcal{O}(c_L^2)$$

Define **log** operator:

$$\mathcal{O}^{\text{log}} = \lim_{\varepsilon \rightarrow 0} \frac{\mathcal{O}^M - \mathcal{O}^L}{\varepsilon}$$

Obtain 2-point correlators:

$$\langle \mathcal{O}^L(z) \mathcal{O}^L(0, 0) \rangle = 0$$

$$\langle \mathcal{O}^L(z) \mathcal{O}^{\text{log}}(0, 0) \rangle = \frac{b_L}{2z^4}$$

$$\langle \mathcal{O}^{\text{log}}(z, \bar{z}) \mathcal{O}^{\text{log}}(0, 0) \rangle = - \frac{b_L \ln(m^2 |z|^2)}{z^4}$$

Critical points and Jordan cells

In terms of leitmotif example:

$$\mathcal{O}^L \sim \psi_1 \quad \mathcal{O}^M \sim \psi_2 \quad \mathcal{O}^{\log} \sim \psi^{\log}$$

Log partner \mathcal{O}^{\log} of EMT has same conformal weights as EMT

Critical points and Jordan cells

In terms of leitmotif example:

$$\mathcal{O}^L \sim \psi_1 \quad \mathcal{O}^M \sim \psi_2 \quad \mathcal{O}^{\log} \sim \psi^{\log}$$

Log partner \mathcal{O}^{\log} of EMT has same conformal weights as EMT

If EMT acquires log partner Hamiltonian cannot be diagonalized

$$H \begin{pmatrix} \mathcal{O}^{\log} \\ \mathcal{O}^L \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} \mathcal{O}^{\log} \\ \mathcal{O}^L \end{pmatrix}$$

Appearance of Jordan cell = defining feature of LCFTs

Note: Jordan cell can be higher rank than 2, but consider only rank 2 case here

LCFTs: Gurarie '93 Reviews on LCFTs: Flohr '01; Gaberdiel '01

LCFT summary

- ▶ $c \rightarrow 0$ limit in CFT intriguing (“ $c = 0$ catastrophe”)

LCFT summary

- ▶ $c \rightarrow 0$ limit in CFT intriguing (“ $c = 0$ catastrophe”)
- ▶ EMT can acquire log partner

LCFT summary

- ▶ $c \rightarrow 0$ limit in CFT intriguing (“ $c = 0$ catastrophe”)
- ▶ EMT can acquire **log partner**
- ▶ Correlators acquire **logarithms**

LCFT summary

- ▶ $c \rightarrow 0$ limit in CFT intriguing (“ $c = 0$ catastrophe”)
- ▶ EMT can acquire **log partner**
- ▶ Correlators acquire **logarithms**
- ▶ Hamiltonian acquires **Jordan cell** structure
- ▶ (**Jordan cells** in quantum mechanics: Berry phases!)

LCFT summary

- ▶ $c \rightarrow 0$ limit in CFT intriguing (“ $c = 0$ catastrophe”)
- ▶ EMT can acquire **log partner**
- ▶ Correlators acquire **logarithms**
- ▶ Hamiltonian acquires **Jordan cell** structure
- ▶ (**Jordan cells** in quantum mechanics: Berry phases!)

Is there a gravity side of the **LCFT** story?



Outline

Gravity in three dimensions

Logarithmic CFTs

$\text{AdS}_3/\text{LCFT}_2$ correspondence

Generalizations

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter
- ▶ Gravity theory should exhibit non-unitarity

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter
- ▶ Gravity theory should exhibit non-unitarity
- ▶ Conformal Ward identities must hold

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter
- ▶ Gravity theory should exhibit non-unitarity
- ▶ Conformal Ward identities must hold
- ▶ Partition function must not be trivial

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter
- ▶ Gravity theory should exhibit non-unitarity
- ▶ Conformal Ward identities must hold
- ▶ Partition function must not be trivial
- ▶ Quasi-normal mode spectrum should reflect appearance of \mathcal{O}^{\log}
- ▶ ...

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter
- ▶ Gravity theory should exhibit non-unitarity
- ▶ Conformal Ward identities must hold
- ▶ Partition function must not be trivial
- ▶ Quasi-normal mode spectrum should reflect appearance of \mathcal{O}^{\log}
- ▶ ...

Cannot be Einstein gravity!

Ingredients for $\text{AdS}_3/\text{LCFT}_2$ correspondence

- ▶ There exists some bulk mode corresponding to the operator \mathcal{O}^{\log}
- ▶ Weights of \mathcal{O}^{\log} must degenerate with weights of \mathcal{O}^L
- ▶ **Jordan cell** structure of $H \sim L_0 + \bar{L}_0$ with respect to \mathcal{O}^{\log} , \mathcal{O}^L
- ▶ **Central charges must be tunable to zero** by some parameter
- ▶ Gravity theory should exhibit non-unitarity
- ▶ Conformal Ward identities must hold
- ▶ Partition function must not be trivial
- ▶ Quasi-normal mode spectrum should reflect appearance of \mathcal{O}^{\log}
- ▶ ...

Cannot be Einstein gravity!

2008-2010:

All items above work for TMG at **critical point** $\mu\ell = 1$!

Critical point in TMG

Central charges in TMG (Kraus & Larsen '05):

$$c_L = \frac{3\ell}{2G} \left(1 - \frac{1}{\mu\ell}\right) \quad c_R = \frac{3\ell}{2G} \left(1 + \frac{1}{\mu\ell}\right)$$

TMG at the critical point is TMG with the tuning

$$\mu\ell = 1$$

between the cosmological constant and the Chern–Simons coupling.
Why special? (Li, Song & Strominger '08)

$$c_L = 0 \quad c_R = \frac{3\ell}{G}$$

Critical point in TMG

Central charges in TMG (Kraus & Larsen '05):

$$c_L = \frac{3\ell}{2G} \left(1 - \frac{1}{\mu\ell}\right) \quad c_R = \frac{3\ell}{2G} \left(1 + \frac{1}{\mu\ell}\right)$$

TMG at the critical point is TMG with the tuning

$$\mu\ell = 1$$

between the cosmological constant and the Chern–Simons coupling.
Why special? (Li, Song & Strominger '08)

$$c_L = 0 \quad c_R = \frac{3\ell}{G}$$

Interesting possibilities:

- ▶ Dual CFT could be chiral (Li, Song & Strominger '08)
- ▶ Dual CFT could be logarithmic (DG & Johansson '08)

Checks of LCFT conjecture

Jordan cell structure

Linearization around AdS background, $g = g^{\text{AdS}} + h$ leads to linearized EOM that are third order PDE

$$G_{\mu\nu}^{(1)} + \frac{1}{\mu} C_{\mu\nu}^{(1)} = (\mathcal{D}^R \mathcal{D}^L \mathcal{D}^M h)_{\mu\nu} = 0 \quad (1)$$

with three mutually commuting first order operators

$$(\mathcal{D}^{L/R})_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} \pm \ell \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha} \quad (\mathcal{D}^M)_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} + \frac{1}{\mu} \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha}$$

Checks of LCFT conjecture

Jordan cell structure

Linearization around AdS background, $g = g^{\text{AdS}} + h$ leads to linearized EOM that are third order PDE

$$G_{\mu\nu}^{(1)} + \frac{1}{\mu} C_{\mu\nu}^{(1)} = (\mathcal{D}^R \mathcal{D}^L \mathcal{D}^M h)_{\mu\nu} = 0 \quad (1)$$

with three mutually commuting first order operators

$$(\mathcal{D}^{L/R})_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} \pm \ell \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha} \quad (\mathcal{D}^M)_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} + \frac{1}{\mu} \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha}$$

Three linearly independent solutions to (1):

$$(\mathcal{D}^L h^L)_{\mu\nu} = 0 \quad (\mathcal{D}^R h^R)_{\mu\nu} = 0 \quad (\mathcal{D}^M h^M)_{\mu\nu} = 0$$

Checks of LCFT conjecture

Jordan cell structure

Linearization around AdS background, $g = g^{\text{AdS}} + h$ leads to linearized EOM that are third order PDE

$$G_{\mu\nu}^{(1)} + \frac{1}{\mu} C_{\mu\nu}^{(1)} = (\mathcal{D}^R \mathcal{D}^L \mathcal{D}^M h)_{\mu\nu} = 0 \quad (1)$$

with three mutually commuting first order operators

$$(\mathcal{D}^{L/R})_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} \pm \ell \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha} \quad (\mathcal{D}^M)_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} + \frac{1}{\mu} \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha}$$

Three linearly independent solutions to (1):

$$(\mathcal{D}^L h^L)_{\mu\nu} = 0 \quad (\mathcal{D}^R h^R)_{\mu\nu} = 0 \quad (\mathcal{D}^M h^M)_{\mu\nu} = 0$$

At **critical point** left (L) and massive (M) branches coincide!

Checks of LCFT conjecture

Jordan cell structure

Linearization around AdS background, $g = g^{\text{AdS}} + h$ leads to linearized EOM that are third order PDE

$$G_{\mu\nu}^{(1)} + \frac{1}{\mu} C_{\mu\nu}^{(1)} = (\mathcal{D}^R \mathcal{D}^L \mathcal{D}^M h)_{\mu\nu} = 0 \quad (1)$$

with three mutually commuting first order operators

$$(\mathcal{D}^{L/R})_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} \pm \ell \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha} \quad (\mathcal{D}^M)_{\mu}{}^{\nu} = \delta_{\mu}^{\nu} + \frac{1}{\mu} \varepsilon_{\mu}^{\alpha\nu} \bar{\nabla}_{\alpha}$$

At **critical point**: get **log** solution (DG & Johansson '08)

$$h_{\mu\nu}^{\log} = \lim_{\mu\ell \rightarrow 1} \frac{h_{\mu\nu}^M(\mu\ell) - h_{\mu\nu}^L}{\mu\ell - 1}$$

with property

$$(\mathcal{D}^L h^{\log})_{\mu\nu} = (\mathcal{D}^M h^{\log})_{\mu\nu} \neq 0, \quad ((\mathcal{D}^L)^2 h^{\log})_{\mu\nu} = 0$$

Log mode exhibits interesting property (DG & Johansson '08):

$$H \begin{pmatrix} h^{\log} \\ h^L \end{pmatrix} = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} h^{\log} \\ h^L \end{pmatrix}$$

Here $H = L_0 + \bar{L}_0 \sim \partial_t$ is the Hamilton operator.

Such a Jordan form of H is defining property of a logarithmic CFT!

Jordan-block structure was main motivation for LCFT conjecture

Checks of LCFT conjecture

Finiteness

Properties of logarithmic mode:

- ▶ Perturbative solution of linearized EOM, but not pure gauge
- ▶ Energy of logarithmic mode is finite

$$E^{\text{log}} = -47/1152 G \ell^3$$

and negative \rightarrow instability (DG & Johansson '08)

- ▶ Logarithmic mode is asymptotically AdS

$$ds^2 = d\rho^2 + \left(\gamma_{ij}^{(0)} e^{2\rho/\ell} + \gamma_{ij}^{(1)} \rho + \gamma_{ij}^{(0)} + \gamma_{ij}^{(2)} e^{-2\rho/\ell} + \dots \right) dx^i dx^j$$

but violates Brown–Henneaux boundary conditions! ($\gamma_{ij}^{(1)}|_{\text{BH}} = 0$)

- ▶ Consistent log boundary conditions replacing Brown–Henneaux (DG & Johansson '08, Martinez, Henneaux & Troncoso '09)
- ▶ Brown–York stress tensor is finite, conserved and traceless, but not chiral (Martinez, Henneaux & Troncoso '09, Maloney, Song & Strominger '09, Ertl, DG & Johansson '09)
- ▶ Log mode persists non-perturbatively, as shown by Hamilton analysis (DG, Jackiw & Johansson '08, Carlip '08)

Checks of LCFT conjecture

Correlators

If LCFT conjecture is correct then following procedure must work:

- ▶ Calculate non-normalizable modes for left, right and logarithmic branches by solving linearized EOM on gravity side
- ▶ According to $\text{AdS}_3/\text{LCFT}_2$ dictionary these non-normalizable modes are sources for corresponding operators in the dual CFT
- ▶ Calculate 2- and 3-point correlators on the gravity side, e.g. by plugging non-normalizable modes into second and third variation of the on-shell action
- ▶ These correlators must coincide with the ones of a LCFT

Checks of LCFT conjecture

Correlators

If LCFT conjecture is correct then following procedure must work:

- ▶ Calculate non-normalizable modes for left, right and logarithmic branches by solving linearized EOM on gravity side
- ▶ According to $\text{AdS}_3/\text{LCFT}_2$ dictionary these non-normalizable modes are sources for corresponding operators in the dual CFT
- ▶ Calculate 2- and 3-point correlators on the gravity side, e.g. by plugging non-normalizable modes into second and third variation of the on-shell action
- ▶ These correlators must coincide with the ones of a LCFT

Except for value of new anomaly b_L no freedom in this procedure. Either it works or it does not work.

Checks of LCFT conjecture

Correlators

If LCFT conjecture is correct then following procedure must work:

- ▶ Calculate non-normalizable modes for left, right and logarithmic branches by solving linearized EOM on gravity side
- ▶ According to $\text{AdS}_3/\text{LCFT}_2$ dictionary these non-normalizable modes are sources for corresponding operators in the dual CFT
- ▶ Calculate 2- and 3-point correlators on the gravity side, e.g. by plugging non-normalizable modes into second and third variation of the on-shell action
- ▶ These correlators must coincide with the ones of a LCFT


Except for value of new anomaly b_L no freedom in this procedure. Either it works or it does not work.

- ▶ Works at level of 2-point correlators (Skenderis, Taylor & van Rees '09, DG & Sachs '09)
- ▶ Works at level of 3-point correlators (DG & Sachs '09)
- ▶ Value of new anomaly: $b_L = -c_R = -3\ell/G$

Checks of LCFT conjecture

1-loop partition function (Gaberdiel, DG & Vassilevich '10)

Structure of low-lying states in LCFT:




Total partition function of Virasoro descendants

$$Z_{\text{LCFT}}^0 = Z_{\Omega} + Z_{\log} = \prod_{n=2}^{\infty} \frac{1}{|1 - q^n|^2} \left(1 + \frac{q^2}{|1 - q|^2} \right)$$

Checks of LCFT conjecture

1-loop partition function (Gaberdiel, DG & Vassilevich '10)

Structure of low-lying states in LCFT:



Total partition function of Virasoro descendants


$$Z_{\text{LCFT}}^0 = Z_{\Omega} + Z_{\log} = \prod_{n=2}^{\infty} \frac{1}{|1 - q^n|^2} \left(1 + \frac{q^2}{|1 - q|^2} \right)$$

Comparison with 1-loop calculation in Euclidean path integral approach to quantum gravity:

Checks of LCFT conjecture

1-loop partition function (Gaberdiel, DG & Vassilevich '10)

Structure of low-lying states in LCFT:



Total partition function of Virasoro descendants

$$Z_{\text{LCFT}}^0 = Z_{\Omega} + Z_{\log} = \prod_{n=2}^{\infty} \frac{1}{|1 - q^n|^2} \left(1 + \frac{q^2}{|1 - q|^2} \right)$$

Comparison with 1-loop calculation in Euclidean path integral approach to quantum gravity:

$$Z_{\text{TMG}} = Z_{\text{LCFT}}^0 + \sum_{h, \bar{h}} N_{h, \bar{h}} q^h \bar{q}^{\bar{h}} \prod_{n=1}^{\infty} \frac{1}{|1 - q^n|^2}$$

All multiplicity coefficients $N_{h, \bar{h}}$ can be shown to be non-negative.

Fairly non-trivial test of the LCFT conjecture!

Checks of LCFT conjecture

Quasi-normal modes (Solodukhin & Sachs '08, Sachs '08)

- ▶ Birmingham, Sachs, Solodukhin '01:
One-to-one correspondence between poles of retarded propagator in CFT and quasi-normal frequencies of linear perturbations of BTZ
- ▶ LCFT should have double pole instead of single pole due to degeneration of operators at critical point
- ▶ Sachs '08:
TMG at critical point has standard right-moving QNM, but no left-moving QNM (pure gauge)
- ▶ Additional QNM from log modes
- ▶ Linear dependence in time of log modes produces the predicted double pole

QNM spectrum compatible with LCFT conjecture:
Double poles in retarded correlators

Outline

Gravity in three dimensions

Logarithmic CFTs

$\text{AdS}_3/\text{LCFT}_2$ correspondence

Generalizations

Summary and generalizations

LCFTs have taught us a great deal about critical TMG!
AdS/LCFT decouples holography from unitarity issues

Summary and generalizations

LCFTs have taught us a great deal about critical TMG!
AdS/LCFT decouples holography from unitarity issues

TMG at the critical point is very special, but to what extent is it generic?

LCFTs have taught us a great deal about critical TMG!
AdS/LCFT decouples holography from unitarity issues

TMG at the critical point is very special, but to what extent is it generic?

- ▶ 3D: generic massive gravity theories (DG, Johansson, Zojer '10)
 - ▶ New massive gravity, generalized massive gravity, ...
 - ▶ higher rank Jordan cells possible in some of these models
 - ▶ qualitatively new LCFTs may arise: no log partner of EMT

LCFTs have taught us a great deal about critical TMG!
AdS/LCFT decouples holography from unitarity issues

TMG at the critical point is very special, but to what extent is it generic?

- ▶ 3D: generic massive gravity theories (DG, Johansson, Zojer '10)
 - ▶ New massive gravity, generalized massive gravity, ...
 - ▶ higher rank Jordan cells possible in some of these models
 - ▶ qualitatively new LCFTs may arise: no log partner of EMT
- ▶ 4D (and higher D): critical gravity (Lü, Pope '11, Deser et. al '11)
2-point correlators match expectations from 3D LCFT (Johansson, Naseh & Zojer '12)

LCFTs have taught us a great deal about critical TMG!
AdS/LCFT decouples holography from unitarity issues

TMG at the critical point is very special, but to what extent is it generic?

- ▶ 3D: generic massive gravity theories (DG, Johansson, Zojer '10)
 - ▶ New massive gravity, generalized massive gravity, ...
 - ▶ higher rank Jordan cells possible in some of these models
 - ▶ qualitatively new LCFTs may arise: no log partner of EMT
- ▶ 4D (and higher D): critical gravity (Lü, Pope '11, Deser et. al '11)
2-point correlators match expectations from 3D LCFT (Johansson, Naseh & Zojer '12)
- ▶ Higher spin log gravity (Chen, Long & Wu '11; Bagchi, Lal, Saha & Sahoo '11)

LCFTs have taught us a great deal about critical TMG!
AdS/LCFT decouples holography from unitarity issues

TMG at the critical point is very special, but to what extent is it generic?

- ▶ 3D: generic massive gravity theories (DG, Johansson, Zojer '10)
 - ▶ New massive gravity, generalized massive gravity, ...
 - ▶ higher rank Jordan cells possible in some of these models
 - ▶ qualitatively new LCFTs may arise: no log partner of EMT
- ▶ 4D (and higher D): critical gravity (Lü, Pope '11, Deser et. al '11)
2-point correlators match expectations from 3D LCFT (Johansson, Naseh & Zojer '12)
- ▶ Higher spin log gravity (Chen, Long & Wu '11; Bagchi, Lal, Saha & Sahoo '11)
- ▶ String construction leading to critical gravity?
Perhaps possible in ABJM context in higgsed phase (Chu & Nilsson '09, Chu, Nastase, Nilsson & Papageorgakis '10)

Thanks to my collaborators ...
... and thanks for your attention!



Vienna group, March 2012



KITP collaboration, May 2012

LCFT: N. Johansson, R. Jackiw, I. Sachs, O. Hohm, M. Gaberdiel, D. Vassilevich,
T. Zojer, S. Ertl, M. Bertin

conformal CS holography: H. Afshar, B. Cvetkovic, N. Johansson, S. Ertl

higher spin holography: M. Gary, R. Rashkov

flat space holography: A. Bagchi, S. Detournay

other holographic aspects: J. Aparicio, E. Lopez, I. Papadimitriou, S. Stricker

Critical points and Jordan cells in quantum mechanics

See “Non-Hermitian quantum mechanics” by Nimrod Moiseyev

Consider the Hamiltonian

$$H = \begin{pmatrix} 1 & \lambda \\ \lambda & -1 \end{pmatrix}$$

with Eigenvalues $E_{\pm} = \pm\sqrt{1 + \lambda^2}$

Critical points and Jordan cells in quantum mechanics

See “Non-Hermitian quantum mechanics” by Nimrod Moiseyev

Consider the Hamiltonian

$$H = \begin{pmatrix} 1 & \lambda \\ \lambda & -1 \end{pmatrix}$$

with Eigenvalues $E_{\pm} = \pm\sqrt{1 + \lambda^2}$

Non-hermitian critical points: $\lambda \rightarrow \pm i$

Eigenvector $c_{\pm} = (\pm i, 1)$ self-orthogonal: $c_{\pm}^2 = 0$

Critical points and Jordan cells in quantum mechanics

See “Non-Hermitian quantum mechanics” by Nimrod Moiseyev

Consider the Hamiltonian

$$H = \begin{pmatrix} 1 & \lambda \\ \lambda & -1 \end{pmatrix}$$

with Eigenvalues $E_{\pm} = \pm\sqrt{1 + \lambda^2}$

Non-hermitian critical points: $\lambda \rightarrow \pm i$

Eigenvector $c_{\pm} = (\pm i, 1)$ self-orthogonal: $c_{\pm}^2 = 0$

Similarity trafo $J = A^{-1}HA$:

$$J = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

Simplest example of Jordan cell in non-hermitian critical quantum mechanics!

Physical significance of critical points?

- ▶ Critical points and Jordan cells require infinite fine-tuning

Physical significance of critical points?

- ▶ Critical points and Jordan cells require infinite fine-tuning
- ▶ Experimentally: can never infinitely fine-tune parameters

Physical significance of critical points?

- ▶ Critical points and Jordan cells require infinite fine-tuning
- ▶ Experimentally: can never infinitely fine-tune parameters
- ▶ Instead: vary parameters adiabatically in vicinity of critical points

$$\text{Eigenvalue: } \lambda = \lambda_c + Re^{i\phi}$$

Physical significance of critical points?

- ▶ Critical points and Jordan cells require infinite fine-tuning
- ▶ Experimentally: can never infinitely fine-tune parameters
- ▶ Instead: vary parameters adiabatically in vicinity of critical points

$$\text{Eigenvalue: } \lambda = \lambda_c + Re^{i\phi}$$

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha \sqrt{R} e^{i\phi/2}$$

Physical significance of critical points?

- ▶ Critical points and Jordan cells require infinite fine-tuning
- ▶ Experimentally: can never infinitely fine-tune parameters
- ▶ Instead: vary parameters adiabatically in vicinity of critical points

$$\text{Eigenvalue: } \lambda = \lambda_c + Re^{i\phi}$$

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha\sqrt{R}e^{i\phi/2}$$

- ▶ Wavefunctions on two branches:

$$|\Psi_{\pm}\rangle = A_{\pm}e^{-i\phi/4}(|\Psi_c\rangle \pm \sqrt{R}e^{i\phi/2}|\chi\rangle)$$

Physical significance of critical points?

- ▶ Critical points and Jordan cells require infinite fine-tuning
- ▶ Experimentally: can never infinitely fine-tune parameters
- ▶ Instead: vary parameters adiabatically in vicinity of critical points

$$\text{Eigenvalue: } \lambda = \lambda_c + Re^{i\phi}$$

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha \sqrt{R} e^{i\phi/2}$$

- ▶ Wavefunctions on two branches:

$$|\Psi_{\pm}\rangle = A_{\pm} e^{-i\phi/4} (|\Psi_c\rangle \pm \sqrt{R} e^{i\phi/2} |\chi\rangle)$$

- ▶ Berry phase: rotate $\phi = 2\pi n$, with $n = 1, 2, 3, 4, \dots$

Berry phase

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha\sqrt{R}e^{i\phi/2}$$

- ▶ Wavefunctions on two branches ($A_- = iA_+$):

$$|\Psi_{\pm}\rangle = A_{\pm}e^{-i\phi/4}(|\Psi_c\rangle \pm \sqrt{R}e^{i\phi/2}|\chi\rangle)$$

- ▶ Berry phase: rotate $\phi = 2\pi n$, with $n = 1, 2, 3, 4, \dots$

- ▶ $n = 1$:

$$E_{\pm}(2\pi) = E_{\mp}(0) \quad |\Psi_{\pm}(2\pi)\rangle = \mp|\Psi_{\mp}(0)\rangle$$

Berry phase

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha\sqrt{R}e^{i\phi/2}$$

- ▶ Wavefunctions on two branches ($A_- = iA_+$):

$$|\Psi_{\pm}\rangle = A_{\pm}e^{-i\phi/4}(|\Psi_c\rangle \pm \sqrt{R}e^{i\phi/2}|\chi\rangle)$$

- ▶ Berry phase: rotate $\phi = 2\pi n$, with $n = 1, 2, 3, 4, \dots$

- ▶ $n = 1$:

$$E_{\pm}(2\pi) = E_{\mp}(0) \quad |\Psi_{\pm}(2\pi)\rangle = \mp|\Psi_{\mp}(0)\rangle$$

- ▶ $n = 2$:

$$E_{\pm}(4\pi) = E_{\pm}(0) \quad |\Psi_{\pm}(4\pi)\rangle = -|\Psi_{\pm}(0)\rangle$$

Berry phase

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha\sqrt{R}e^{i\phi/2}$$

- ▶ Wavefunctions on two branches ($A_- = iA_+$):

$$|\Psi_{\pm}\rangle = A_{\pm}e^{-i\phi/4}(|\Psi_c\rangle \pm \sqrt{R}e^{i\phi/2}|\chi\rangle)$$

- ▶ Berry phase: rotate $\phi = 2\pi n$, with $n = 1, 2, 3, 4, \dots$

- ▶ $n = 1$:

$$E_{\pm}(2\pi) = E_{\mp}(0) \quad |\Psi_{\pm}(2\pi)\rangle = \mp|\Psi_{\mp}(0)\rangle$$

- ▶ $n = 2$:

$$E_{\pm}(4\pi) = E_{\pm}(0) \quad |\Psi_{\pm}(4\pi)\rangle = -|\Psi_{\pm}(0)\rangle$$

- ▶ $n = 4$:

$$E_{\pm}(8\pi) = E_{\pm}(0) \quad |\Psi_{\pm}(8\pi)\rangle = +|\Psi_{\pm}(0)\rangle$$

Berry phase

- ▶ Energies on two branches:

$$E_{\pm}(\phi) = E_c \pm \alpha\sqrt{R}e^{i\phi/2}$$

- ▶ Wavefunctions on two branches ($A_- = iA_+$):

$$|\Psi_{\pm}\rangle = A_{\pm}e^{-i\phi/4}(|\Psi_c\rangle \pm \sqrt{R}e^{i\phi/2}|\chi\rangle)$$

- ▶ Berry phase: rotate $\phi = 2\pi n$, with $n = 1, 2, 3, 4, \dots$

- ▶ $n = 1$:

$$E_{\pm}(2\pi) = E_{\mp}(0) \quad |\Psi_{\pm}(2\pi)\rangle = \mp|\Psi_{\mp}(0)\rangle$$

- ▶ $n = 2$:

$$E_{\pm}(4\pi) = E_{\pm}(0) \quad |\Psi_{\pm}(4\pi)\rangle = -|\Psi_{\pm}(0)\rangle$$

- ▶ $n = 4$:

$$E_{\pm}(8\pi) = E_{\pm}(0) \quad |\Psi_{\pm}(8\pi)\rangle = +|\Psi_{\pm}(0)\rangle$$

Physical significance of critical points: geometrical (Berry) phases!

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation
- ▶ For sufficient amount of disorder perturbation theory breaks down — random critical point

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation
- ▶ For sufficient amount of disorder perturbation theory breaks down — random critical point
- ▶ Infamous denominator in correlators:

$$\overline{\langle \mathcal{O}(z) \mathcal{O}(0) \rangle} = \int \mathcal{D}V P[V] \frac{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right) \mathcal{O}(z) \mathcal{O}(0)}{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right)}$$

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation
- ▶ For sufficient amount of disorder perturbation theory breaks down — random critical point
- ▶ Infamous denominator in correlators:

$$\overline{\langle \mathcal{O}(z) \mathcal{O}(0) \rangle} = \int \mathcal{D}V P[V] \frac{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right) \mathcal{O}(z) \mathcal{O}(0)}{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right)}$$

- ▶ Different ways to deal with denominator (replica trick, SUSY)

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation
- ▶ For sufficient amount of disorder perturbation theory breaks down — random critical point
- ▶ Infamous denominator in correlators:

$$\overline{\langle \mathcal{O}(z) \mathcal{O}(0) \rangle} = \int \mathcal{D}V P[V] \frac{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right) \mathcal{O}(z) \mathcal{O}(0)}{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right)}$$

- ▶ Different ways to deal with denominator (replica trick, SUSY)
- ▶ Result: operators degenerate and correlators acquire logarithmic behavior, exactly as in LCFT (Cardy '99)

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation
- ▶ For sufficient amount of disorder perturbation theory breaks down — random critical point
- ▶ Infamous denominator in correlators:

$$\overline{\langle \mathcal{O}(z) \mathcal{O}(0) \rangle} = \int \mathcal{D}V P[V] \frac{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right) \mathcal{O}(z) \mathcal{O}(0)}{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right)}$$

- ▶ Different ways to deal with denominator (replica trick, SUSY)
- ▶ Result: operators degenerate and correlators acquire logarithmic behavior, exactly as in LCFT (Cardy '99)
- ▶ Exploit LCFTs to compute correlators of quenched random systems

Systems with quenched disorder

LCFTs arise in systems with quenched disorder.

- ▶ Quenched disorder: systems with random variable that does not evolve in time
- ▶ Examples: spin glasses, quenched random magnets, dilute self-avoiding polymers, percolation
- ▶ For sufficient amount of disorder perturbation theory breaks down — random critical point
- ▶ Infamous denominator in correlators:

$$\overline{\langle \mathcal{O}(z) \mathcal{O}(0) \rangle} = \int \mathcal{D}V P[V] \frac{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right) \mathcal{O}(z) \mathcal{O}(0)}{\int \mathcal{D}\phi \exp \left(- I[\phi] - \int d^2 z' V(z') \mathcal{O}(z') \right)}$$

- ▶ Different ways to deal with denominator (replica trick, SUSY)
- ▶ Result: operators degenerate and correlators acquire logarithmic behavior, exactly as in LCFT (Cardy '99)
- ▶ Exploit LCFTs to compute correlators of quenched random systems
- ▶ Idea: **Apply $\text{AdS}_3/\text{LCFT}_2$ to describe strongly coupled LCFTs!**

Some literature on condensed matter applications of LCFTs

- ▶ Cardy '99 Logarithmic correlations in Quenched Random Magnets and Polymers
- ▶ Gurarie & Ludwig '99 Conformal algebras of 2D disordered systems
- ▶ Rahimi Tabar '00 Quenched Averaged Correlation Functions of the Random Magnets
- ▶ Reviews: Flohr '01 and Gaberdiel '01

Some literature on condensed matter applications of LCFTs

- ▶ Cardy '99 Logarithmic correlations in Quenched Random Magnets and Polymers
- ▶ Gurarie & Ludwig '99 Conformal algebras of 2D disordered systems
- ▶ Rahimi Tabar '00 Quenched Averaged Correlation Functions of the Random Magnets
- ▶ Reviews: Flohr '01 and Gaberdiel '01

More applications awaiting to be discovered!

