

Recent developments in 2d (0,2) theories

Eric Sharpe
Virginia Tech

C. Closset, W. Gu, B. Jia, ES, arXiv:1512.08058
J. Guo, V. Lu, ES, arXiv:1512.08586 & 1605.01410
Z. Chen, ES, R. Wu, arXiv: 1603.09634

Over the last decade, there's been a *tremendous* amount of progress in perturbative string compactifications.

A few of my favorite examples:

- nonpert' realizations of geometry (Pfaffians, double covers)
(Hori-Tong '06, Caldararu et al '07,...)
- perturbative GLSM's for Pfaffians (Hori '11, Jockers et al '12,...)
- non-birational GLSM phases - physical realization of homological projective duality
(Hori-Tong '06, Caldararu et al '07, Ballard et al '12; Kuznetsov '05-'06,...)
- examples of closed strings on noncommutative res'ns
(Caldararu et al '07, Addington et al '12, ES '13)
- localization techniques: new GW & elliptic genus computations, role of Gamma classes, ...
(Benini-Cremonesi '12, Doroud et al '12; Jockers et al '12, Halverson et al '13, Hori-Romo '13, Benini et al '13,)
- heterotic strings: nonpert' corrections, 2d dualities/trialities, non-Kahler moduli (many)

I'll focus today on just one....

Today I'll restrict to

- heterotic strings: nonpert' corrections, 2d dualities/trialities, non-Kahler moduli

My goal today: an overview of progress towards computing worldsheet instanton effects, one of the outstanding problems in perturbative heterotic string compactifications.

Briefly, we need to generalize instanton corrections and mirror symmetry to heterotic theories, and some progress has been made.

Results include quantum sheaf cohomology, a generalization of quantum cohomology.

Review gen'l aspects next....

Some background.

In 10d, a heterotic string describes metric & gauge field.

To compactify, must specify not only a space X , but also a holomorphic vector bundle \mathcal{E} on that space, satisfying consistency conditions

$$[\text{tr } F \wedge F] = [\text{tr } R \wedge R] \quad \text{ch}_2(\mathcal{E}) = \text{ch}_2(TX)$$

Described on worldsheet by 2d (0,2) susy theory.

Simplest case: $\mathcal{E} = TX$, corresponding to (2,2) susy.

“embed the spin connection in gauge connection”

Simplest case: compactification on a Calabi-Yau with
gauge bundle $\mathcal{E} =$ tangent bundle TX ,
(`embedding the spin connection' = (2,2) locus)

In this case, we know basics:

- massless states (inc. moduli)
— counted by cohomology of the CY; `chiral ring'
- Yukawa couplings, superpotentials
(inc. nonperturbative corrections)

Nonperturbative corrections = GW inv'ts

$\overline{27}^3$ = A model TFT computation
OPE's => quantum cohomology

27^3 = B model TFT computation

More gen'l case: compactification on a Calabi-Yau with gauge bundle $\mathcal{E} \neq$ tangent bundle TX .

(Worldsheet has (0,2) susy.)

- massless states (inc. moduli)
- counted by bundle-valued forms on the CY
(= sheaf cohomology, eg, $\overline{\mathbf{27}} \sim H^1(X, \mathcal{E}^*)$)
- Yukawa couplings, superpotentials
(inc. nonperturbative corrections)

Nonperturbative corrections \neq GW inv'ts

$\overline{\mathbf{27}}^3 = A/2$ model computation

OPE's \Rightarrow quantum sheaf cohomology

$\mathbf{27}^3 = B/2$ model computation

- Yukawa couplings, superpotentials
(inc. nonperturbative corrections)


Nonperturbative corrections \neq GW inv'ts

$$\overline{\mathbf{27}}^3 = A/2 \text{ model computation}$$

$$\mathbf{27}^3 = B/2 \text{ model computation}$$

But we know far less for (0,2) than for (2,2)!

Understanding these nonperturbative corrections is a central issue in perturbative heterotic strings on CYs.



quantum sheaf cohomology
(0,2) mirror symmetry

generalizing ordinary quantum cohomology & mirror symmetry.

In the next few minutes before the break,
I'll outline in more detail

- correspondence of massless states to sheaf cohomology
- form of quantum corrections to correlation functions
- basic aspects of the $A/2$, $B/2$ analogues of TFT's
- examples of results for quantum sheaf cohomology rings

Then, after the break, I'll walk through details of computations.

First, before outlining Yukawa couplings,
how to count massless states?

Let X be Calabi-Yau 3-fold, for simplicity.

Let \mathcal{E} denote the gauge bundle.

(Distler-Greene '88)

Rank 3 bundle: low-energy $E_6 \times E_8$

$$\overline{\mathbf{27}} \sim H^1(X, \mathcal{E}^*)$$

$$\mathbf{27} \sim H^1(X, \mathcal{E})$$

$$(2,2): \quad \sim H^1(X, T^*X) \text{ Kähler} \quad \sim H^1(X, TX) \text{ complex}$$

Rank 4 bundle: low-energy $\text{Spin}(10) \times E_8$

$$\mathbf{16} \sim H^1(X, \mathcal{E})$$

$$\mathbf{10} \sim H^1(X, \wedge^2 \mathcal{E})$$

Rank 5 bundle: low-energy $SU(5) \times E_8$

$$\mathbf{10} \sim H^1(X, \mathcal{E})$$

$$\overline{\mathbf{5}} \sim H^1(X, \wedge^2 \mathcal{E})$$

What are their couplings?

What are their couplings?

Suppose bundle is rank 3, for simplicity,
so that we have low-energy $E_6 \times E_8$.

$$\overline{27}^3 = \int_X \omega_1 \wedge \omega_2 \wedge \omega_3 + \mathcal{O}(q) \quad \text{where } \omega_i \in H^1(X, \mathcal{E}^*)$$

No perturbative loop corrections, but there are nonperturbative corrections.

(Dine-Seiberg-Wen-Witten '86)

Example: (2,2) quintic

(Candelas, de la Ossa,
Green, Parkes, '91)

$$\overline{27}^3 = 5 + \sum_{k=1}^{\infty} \frac{n_k k^3 q^k}{1 - q^k} = 5 + 2875q + 4876875 q^2 + \dots$$

intersection number

(Strominger '85)

nonperturbative
contributions

n_k = Gromov-Witten
invariants

How to compute nonperturbative corrections in (0,2) cases?

How to compute nonperturbative corrections in $(0,2)$ cases?

Historically, on the $(2,2)$ locus,
used mirror symmetry.

For $(0,2)$, would need a generalization called
 $(0,2)$ mirror symmetry.

Some results do exist — state of the art is a version of
Batyrev's mirror map due to **Melnikov-Plesser '10** — but we
have not yet worked out analogue of flat coordinates or how to
compute nonperturbative corrections using $(0,2)$ mirrors alone.

We'll do this directly instead....

How to compute nonperturbative corrections in (0,2) cases?

It's convenient to work in an analogue of a TFT.

On (2,2) locus,

$$\overline{\mathbf{27}}^3 = \langle V_f^{16} V_b^{10} V_f^{16} \rangle_{\text{phys}} = \langle V^3 \rangle_{\text{A TFT}}$$

$$\mathbf{27}^3 = \langle V_f^{16} V_b^{10} V_f^{16} \rangle_{\text{phys}} = \langle V^3 \rangle_{\text{B TFT}}$$

& the TFT expressions are convenient for computations.

There are analogues for more general (0,2) theories; these are A/2, B/2 pseudo-TFT's, which also have the property

$$\overline{\mathbf{27}}^3 = \langle V_f^{16} V_b^{10} V_f^{16} \rangle_{\text{phys}} = \langle V^3 \rangle_{\text{A/2 TFT}}$$

$$\mathbf{27}^3 = \langle V_f^{16} V_b^{10} V_f^{16} \rangle_{\text{phys}} = \langle V^3 \rangle_{\text{B/2 TFT}}$$

The $A/2$, $B/2$ pseudo-TFT's

These $(0,2)$ NLSM's have two anomalous global $U(1)$'s:

- a right-moving $U(1)_R$
- a canonical left-moving $U(1)$,
rotating the phase of all left fermions,
which becomes $U(1)_L$ on $(2,2)$ locus

If $\det \mathcal{E}^{\pm 1} \cong K_X$, then a nonanomalous $U(1)$ exists along which we can twist right & left moving fermions.

There are two distinct possibilities,
which on $(2,2)$ locus become the A , B model TFT's,
and are called the $A/2$, $B/2$ models.

A little more explicitly:

(0,2) NLSM has Lagrangian density

$$\mathcal{L} = g_{i\bar{j}} \partial \phi^{\bar{j}} \bar{\partial} \phi^i + i g_{i\bar{j}} \psi_{+}^{\bar{j}} D_{-} \psi_{+}^i + i h_{a\bar{b}} \lambda_{-}^{\bar{b}} D_{+} \lambda_{-}^a + F_{i\bar{j}a\bar{b}} \psi_{+}^i \psi_{+}^{\bar{j}} \lambda_{-}^a \lambda_{-}^{\bar{b}}$$
$$\psi_{+} \sim TX \quad \lambda_{-} \sim \mathcal{E}$$

subject to Green-Schwarz condition: $\text{ch}_2(TX) = \text{ch}_2(\mathcal{E})$

A/2 twist: take $\psi_{+}^i, \lambda_{-}^{\bar{a}}$ to be scalars

B/2 twist: take $\psi_{+}^{\bar{i}}, \lambda_{-}^{\bar{a}}$ to be scalars

so we get a scalar half of susy — but this BRST operator is purely right-moving, so this not a standard TFT.

In order for this twist to be anomaly-free, there are constraints..

A/2 model: Exists when $(\det \mathcal{E})^{-1} \cong K_X$

(on (2,2) locus, always possible; reduces to A model)

States: $H^\bullet(X, \wedge^\bullet \mathcal{E}^*)$

B/2 model: Exists when $\det \mathcal{E} \cong K_X$

(on (2,2) locus, requires $K_X^{\otimes 2} \cong \mathcal{O}_X$; reduces to B model)

States: $H^\bullet(X, \wedge^\bullet \mathcal{E})$

Exchanging $\mathcal{E} \leftrightarrow \mathcal{E}^*$ swaps the A/2, B/2 models.

(Physically, just a complex conjugation of left movers.)

What do the A, A/2 model correlation functions look like?

Classical contributions, schematically:

A model: Classical contribution:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \int_X \omega_1 \wedge \cdots \wedge \omega_n = \int_X (\text{top form})$$

$\omega_i \in H^{p_i, q_i}(X)$

A/2 model: Classical contribution:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \int_X \omega_1 \wedge \cdots \wedge \omega_n \quad \omega_i \in H^{q_i}(X, \wedge^{p_i} \mathcal{E}^*)$$

Now, $\omega_1 \wedge \cdots \wedge \omega_n \in H^{\text{top}}(X, \wedge^{\text{top}} \mathcal{E}^*) = H^{\text{top}}(X, K_X)$

using the anomaly constraint $\det \mathcal{E}^* \cong K_X$

Again, a top form, so get a number.

What do the A, A/2 model correlation functions look like?

Instanton sectors, schematically:

A model:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \int_{\mathcal{M}} \omega_1 \wedge \cdots \wedge \omega_n = \int_{\mathcal{M}} (\text{top - form})$$

$\omega_i \in H^{p_i, q_i}(\mathcal{M})$

where \mathcal{M} is moduli space of worldsheet instantons.

A/2 model:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \int_{\mathcal{M}} \omega_1 \wedge \cdots \wedge \omega_n \quad \omega_i \in H^{q_i}(\mathcal{M}, \wedge^{p_i} \mathcal{F}^*)$$

where \mathcal{F} is sheaf on \mathcal{M} induced by \mathcal{E} .

Now, $\omega_1 \wedge \cdots \wedge \omega_n \in H^{\text{top}}(\mathcal{M}, \wedge^{\text{top}} \mathcal{F}^*)$

so need to explain how to get top-form etc....

What do the A , $A/2$ model correlation functions look like?

To actually define A model correlation functions,
need to compactify \mathcal{M} .

To actually define $A/2$ model correlation functions,
need to not only compactify \mathcal{M} ,
but also extend \mathcal{F} over compactification divisor,
consistent with symmetries.

Then, formally, get a top-form so long as no anomalies:

$$\left. \begin{array}{l} \wedge^{\text{top}} \mathcal{E}^* \cong K_X \\ \text{ch}_2(\mathcal{E}) = \text{ch}_2(TX) \end{array} \right\} \xrightarrow{\text{GRR}} \wedge^{\text{top}} \mathcal{F}^* \cong K_{\mathcal{M}}$$

All of this has been done.

(Katz-ES [hep-th/0406226](#), ...,
[Donagi-Guffin-Katz-ES 1110.3751](#), [.3752](#))

Resulting corr' f'ns encoded in quantum (sheaf) cohomology

Mathematically:

Quantum sheaf cohomology refers to quantum-corrected product structure on

$$\bigoplus H^\bullet(X, \wedge^\bullet \mathcal{E}^*)$$

where

$$\text{ch}_2(\mathcal{E}) = \text{ch}_2(TX) \quad \text{and} \quad \det \mathcal{E}^* \cong K_X$$

and the classical product structure is of form

$$H^{i_1}(X, \wedge^{j_1} \mathcal{E}^*) \otimes H^{i_2}(X, \wedge^{j_2} \mathcal{E}^*) \longrightarrow H^{i_1+i_2}(X, \wedge^{j_1+j_2} \mathcal{E}^*)$$

Reduces to ordinary quantum cohomology when $\mathcal{E} = TX$

For example: $\bigoplus H^\bullet(X, \wedge^\bullet \mathcal{E}^*) \rightsquigarrow \bigoplus H^\bullet(X, \Omega^\bullet) = \bigoplus H^{\bullet, \bullet}(X)$

(In physics, charged matter states in the heterotic compactification are counted by these sheaf cohom' gps.)

Examples of resulting mathematical structures:

$$\mathbb{P}^1 \times \mathbb{P}^1 :$$

Ordinary quantum cohomology ring is

$$\mathbb{C}[\sigma, \tilde{\sigma}] / (\sigma^2 = q, \tilde{\sigma}^2 = \tilde{q})$$

For \mathcal{E} a deformation of the tangent bundle defined by
four 2x2 matrices A, B, C, D,

quantum sheaf cohomology ring is

$$\mathbb{C}[\sigma, \tilde{\sigma}] / (\det(A\sigma + B\tilde{\sigma}) = q, \det(C\sigma + D\tilde{\sigma}) = \tilde{q})$$

Recover tangent bundle & ordinary quantum cohomology
when $A=D=I, B=C=0$.

We'll see details later.

G(k,n):

Ordinary quantum cohomology ring is

$$\mathbb{C}[\sigma_{(1)}, \dots, \sigma_{(n-k)}] / \langle D_{k+1}, \dots, D_{n-1}, D_n + (-)^n q \rangle =$$

$$\mathbb{C}[\sigma_{(1)}, \sigma_{(2)}, \dots] / \langle D_{k+1}, D_{k+2}, \dots, \sigma_{(n-k+1)}, \dots, \sigma_{(n-1)},$$

$$\sigma_{(n)} + q, \sigma_{(n+1)} + q\sigma_{(1)}, \dots \rangle$$

where $D_m = \det (\sigma_{(1+j-i)})_{1 \leq i, j \leq m}$

For a def' of tangent bundle, the quantum sheaf coh' ring is

$$\mathbb{C}[\sigma_{(1)}, \sigma_{(2)}, \dots,] / \langle D_{k+1}, D_{k+2}, \dots, R_{(n-k+1)}, \dots, R_{(n-1)},$$

$$R_{(n)} + q, R_{(n+1)} + q\sigma_{(1)}, R_{(n+2)} + q\sigma_{(2)}, \dots \rangle$$

where $R_{(r)} = \sum_{i=0}^{\min(r,n)} I_i \sigma_{(r-i)} \sigma_{(1)}^i$ for I_i the coeff's of char' poly' of matrix defining deformation.

We'll see details later.

After the break:

- Susy localization in $A/2$ model for def's of $(2,2)$ theories
- Examples: $\mathbb{P}^1 \times \mathbb{P}^1$, \mathbb{F}_n , $G(k, n)$
 - new expressions for old results: JKG residues
 - new results: nonabelian GLSM's
- $(0,2)$ Toda duals
- Analogous computations in dual $B/2$ theories

Break

Outline of the rest of this talk:

- Susy localization in $A/2$ model for def's of $(2,2)$ theories
- Examples: $\mathbb{P}^1 \times \mathbb{P}^1$, \mathbb{F}_n , $G(k, n)$
 - new expressions for old results: JKG residues
 - new results: nonabelian GLSM's
- $(0,2)$ Toda duals
- Analogous computations in dual $B/2$ theories

Susy localization

I'll first discuss $A/2$ theories obtained by deforming off the $(2,2)$ locus, generalizing A model susy localization described in

[Benini-Zaffaroni 1504.03698](#)

[Closset-Cremonesi-Park 1504.06308](#)

Corresponding $(0,2)$ GLSM's will have a Coulomb branch, along which we shall work.

Schematically, correlation functions take general form

$$\langle f(\sigma) \rangle = \sum_{m \in \mathbb{Z}} \text{JKG} - \text{Res}_{\sigma=0} \left\{ Z^{1-\text{loop}} q^m f(\sigma) \right\}$$

$$\text{for } Z^{1-\text{loop}} = \frac{\det \mathcal{O}_{\text{fermi}}}{\det \mathcal{O}_{\text{bose}}}$$

Susy localization

$$Z^{1\text{-loop}} = \frac{\det \mathcal{O}_{\text{fermi}}}{\det \mathcal{O}_{\text{bose}}}$$

For deformations off the (2,2) locus, in a GLSM,
 ψ_+, ψ_- have same gauge charges.

Fermi interactions: $\bar{\psi}_-^i \psi_+^j E_i^j + \bar{\psi}_+^j \psi_-^i (E_i^j)^*$

$$\mathcal{O}_{\text{fermi}} = \begin{bmatrix} E_1^1 & D_+ & E_1^2 & 0 & \cdots \\ D_- & (E_1^1)^* & 0 & (E_2^1)^* & \cdots \\ E_2^1 & 0 & E_2^2 & D_+ & \cdots \\ 0 & (E_1^2)^* & D_- & (E_2^2)^* & \cdots \\ \vdots & & & & \ddots \end{bmatrix}$$

$$\det \mathcal{O}_{\text{fermi}} = (S(\det E))^{|b+1|} \prod_{n \geq 1} \left[\sum_{k=0}^N t_n^{2k} \left(\sum_{i_1 < i_2 < \cdots < i_k, j_1 < j_2 < \cdots < j_k} \left| \tilde{E}_{i_1 \cdots i_k j_1 \cdots j_k} \right|^2 \right) \right]^{2n+|b+1|}$$

where $t_n = n(n + |b + 1|)$ $b = Q(\mathbf{m})$

Susy localization

$$Z^{1\text{-loop}} = \frac{\det \mathcal{O}_{\text{fermi}}}{\det \mathcal{O}_{\text{bose}}}$$

Bosonic potential:

$$|E^i(\phi)|^2 = \sum_i \left(\sum_j |E_j^i|^2 |\phi^j|^2 \right) + \sum_{i \neq j} \left(\sum_k (E_i^k)^* E_j^k \right) \bar{\phi}^i \phi^j$$

$$\mathcal{O}_{\text{bose}} = \begin{bmatrix} -D^2 + |E_1^1|^2 + \dots + |E_1^N|^2 & (E_1^1)^* E_2^1 + \dots + (E_1^N)^* E_2^N & \dots \\ E_1^1 (E_2^1)^* + \dots + E_1^N (E_2^N)^* & -D^2 + |E_2^1|^2 + \dots + |E_2^N|^2 & \dots \\ \vdots & & \ddots \end{bmatrix}$$

$$\det \mathcal{O}_{\text{bose}} = \prod_{n \geq 0} \left[\sum_{k=0}^N t_n^{2k} \left(\sum_{i_1 < i_2 < \dots < i_k, j_1 < j_2 < \dots < j_k} \left| \tilde{E}_{i_1 \dots i_k j_1 \dots j_k} \right|^2 \right) \right]^{2n+|b|+1}$$

where $t_n = \frac{1}{2} (2n(n+1) + (2n+1)|b| - b)$

Susy localization

Putting this together, can show

$$Z^{1-\text{loop}} = \frac{\det \mathcal{O}_{\text{fermi}}}{\det \mathcal{O}_{\text{bose}}} = \left(\frac{1}{\det E} \right)^{Q(\mathfrak{m})+1}$$

so *schematically* correlation functions take form

$$\begin{aligned} \langle f(\sigma) \rangle &= \sum_{\mathfrak{m} \in \mathbb{Z}} \text{JKG} - \text{Res}_{\sigma=0} \left\{ Z^{1-\text{loop}} q^{\mathfrak{m}} f(\sigma) \right\} \\ &= \sum_{\mathfrak{m} \in \mathbb{Z}} \text{JKG} - \text{Res}_{\sigma=0} \left\{ \left(\frac{1}{\det E} \right)^{Q(\mathfrak{m})+1} q^{\mathfrak{m}} f(\sigma) \right\} \end{aligned}$$

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

Build a (0,2) theory that deforms (2,2) model.

Math: $0 \longrightarrow \mathcal{O}^2 \xrightarrow{*} \mathcal{O}(1,0)^2 \oplus \mathcal{O}(0,1)^2 \longrightarrow \mathcal{E} \longrightarrow 0$

$$* = \begin{bmatrix} Ax & Bx \\ C\tilde{x} & D\tilde{x} \end{bmatrix}$$

x, \tilde{x} vectors of homogeneous coordinates,

A, B, C, D 2×2 matrices describing deformation

(2,2) locus: $A = D = I_{2 \times 2}, \quad B = C = 0$

Physics....

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

Build a (0,2) theory that deforms (2,2) model.

Physics:

x^i, \tilde{x}^i chiral superfields charge (1,0), (0,1)

$\Lambda^i, \tilde{\Lambda}^i$ Fermi superfields charge (1,0), (0,1) s.t.

$$\bar{D}_+ \Lambda^i = A_j^i \sigma x^j + B_j^i \tilde{\sigma} x^j \quad \bar{D}_+ \tilde{\Lambda}^i = C_j^i \sigma \tilde{x}^j + D_j^i \tilde{\sigma} \tilde{x}^j$$

σ neutral (adj-valued) chiral superfield

no superpotential

On (2,2) locus, x^i, Λ^i combine into (2,2) chiral superfields,
 $\tilde{x}^i, \tilde{\Lambda}^i$ combine into (2,2) chiral superfields, and
 σ part of (2,2) vector multiplet.

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

Localization computation: (genus zero)

$$\langle f(\sigma, \tilde{\sigma}) \rangle = \sum_{m_1, m_2 \in \mathbb{Z}} \text{JKG} - \text{Res}_{\sigma=\tilde{\sigma}=0} \left\{ \left(\frac{1}{\det E} \right)^{m_1+1} \left(\frac{1}{\det \tilde{E}} \right)^{m_2+1} q^{m_1} \tilde{q}^{m_2} f(\sigma, \tilde{\sigma}) \right\}$$

$$\text{for } E = A\sigma + B\tilde{\sigma}, \quad \tilde{E} = C\sigma + D\tilde{\sigma}$$

Note: Looks like a TFT result — no propagators, no worldsheet position dependence — but this is not quite TFT.

“Non-topological TFT”

How can that be?

“Non-topological TFT”

The basic reason we're getting a TFT-like structure, albeit not in an actual TFT, is that the OPE's close on dim zero $A/2$ op's.

(Adams-Distler-Ernebjerg '05) argued that e.g. in an open patch on moduli space containing $(2,2)$ locus, the OPE's of the $A/2$ model operators close into other $A/2$ model operators.

For conformal cases, combination of

- right-moving $N=2$ algebra to bound dimensions
 - worldsheet conformal invariance to relate left, right dim's
- to argue closure on patches.

Since operators have dim' zero, & OPE's close, no worldsheet dependence in correlation functions.

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

Let's take another look at the result:

$$\langle f(\sigma, \tilde{\sigma}) \rangle = \sum_{m_1, m_2 \in \mathbb{Z}} \text{JKG} - \text{Res}_{\sigma=\tilde{\sigma}=0} \left\{ \left(\frac{1}{\det E} \right)^{m_1+1} \left(\frac{1}{\det \tilde{E}} \right)^{m_2+1} q^{m_1} \tilde{q}^{m_2} f(\sigma, \tilde{\sigma}) \right\}$$

for $E = A\sigma + B\tilde{\sigma}, \quad \tilde{E} = C\sigma + D\tilde{\sigma}$

Inserting a factor of, say, $\det E$ in the correlation f'n
is equivalent to shifting q .

Quantum sheaf cohomology ring rel'ns:

$$\det E = q, \quad \det \tilde{E} = \tilde{q}$$

This result already known (for all toric varieties w/ def's):

Physics: McOrist-Melnikov 0810.0012

Math: Donagi-Guffin-Katz-ES 1110.3751, .3752

but the derivation is new.

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

Compare

Quantum sheaf cohomology (q.s.c.) ring rel'ns:

$$\det(A\sigma + B\tilde{\sigma}) = q, \quad \det(C\sigma + D\tilde{\sigma}) = \tilde{q}$$

Ordinary quantum cohomology ring rel'ns:

$$\sigma^2 = q, \quad \tilde{\sigma}^2 = \tilde{q}$$

On the (2,2) locus, where $A = D = I_{2 \times 2}$, $B = C = 0$

quantum sheaf cohomology reduces to
ordinary quantum cohomology.

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

2-pt correlation functions:

$$\langle \sigma \sigma \rangle = -\alpha^{-1} \Gamma_1 \quad \langle \sigma \tilde{\sigma} \rangle = \alpha^{-1} \Delta \quad \langle \tilde{\sigma} \tilde{\sigma} \rangle = -\alpha^{-1} \Gamma_2$$

where

$$\Gamma_1 = \gamma_{AB} \det D - \gamma_{CD} \det B \quad \Gamma_2 = \gamma_{CD} \det A - \gamma_{AB} \det C$$

$$\gamma_{AB} = \det(A + B) - \det A - \det B$$

$$\gamma_{CD} = \det(C + D) - \det C - \det D$$

$$\Delta = \det A \det D - \det B \det C$$

$$\alpha = \Delta^2 - \Gamma_1 \Gamma_2$$

$\{\alpha = 0\}$ = locus where bundle degenerates

JKG residue results match Cech cohomology computation.

Example: $\mathbb{P}^1 \times \mathbb{P}^1$

2-pt correlation functions:

$$\langle \sigma \sigma \rangle = -\alpha^{-1} \Gamma_1 \quad \langle \sigma \tilde{\sigma} \rangle = \alpha^{-1} \Delta \quad \langle \tilde{\sigma} \tilde{\sigma} \rangle = -\alpha^{-1} \Gamma_2$$

Can show these 2-pt functions obey

$$\langle \det(A\sigma + B\tilde{\sigma}) \rangle = 0 \quad \langle \det(C\sigma + D\tilde{\sigma}) \rangle = 0$$

matching classical limit of q.s.c. relations.

Can also compute higher-pt functions.

They also match Cech computations, and obey suitable OPE's;
for brevity, let's move on.

Example: Hirzebruch surfaces \mathbb{F}_n

Build a (0,2) theory that deforms (2,2) model.

Math:

$$0 \longrightarrow \mathcal{O}^2 \xrightarrow{*} \mathcal{O}(1,0)^2 \oplus \mathcal{O}(n,1) \oplus \mathcal{O}(0,1) \longrightarrow \mathcal{E} \longrightarrow 0$$

$$* = \begin{bmatrix} Ax & Bx \\ \gamma_1 w + s f_n(x_1, x_2) & \beta_1 w + s g_n(x_1, x_2) \\ \gamma_2 s & \beta_2 s \end{bmatrix}$$

Physics:

$$\overline{D}_+ \Lambda^i = A_j^i \sigma x^j + B_j^i \tilde{\sigma} x^j$$

$$\overline{D}_+ \Lambda_w = \sigma(\gamma_1 w + s f_n) + \tilde{\sigma}(\beta_1 w + s g_n)$$

$$\overline{D}_+ \Lambda_s = \sigma \gamma_2 s + \tilde{\sigma} \beta_2 s$$

— depends upon deg n polynomials f_n, g_n ;
however, they don't contribute to correlation functions:

Example: Hirzebruch surfaces \mathbb{F}_n

Localization computation:

$$\langle f(\sigma, \tilde{\sigma}) \rangle = \sum_{m_1, m_2 \in \mathbb{Z}} \text{JKG} - \text{Res}_{\sigma=\tilde{\sigma}=0} \left\{ \left(\frac{1}{\det E} \right)^{m_1+1} \left(\frac{1}{Q_w} \right)^{nm_1+m_2+1} \left(\frac{1}{Q_s} \right)^{m_2+1} q^{m_1} \tilde{q}^{m_2} f(\sigma, \tilde{\sigma}) \right\}$$

$$E = A\sigma + B\tilde{\sigma}$$

$$Q_w = \gamma_1\sigma + \beta_1\tilde{\sigma}$$

$$Q_s = \gamma_2\sigma + \beta_2\tilde{\sigma}$$

Can read off quantum sheaf cohomology ring rel'ns:

$$(\det E)Q_w^n = q \quad Q_s Q_w = \tilde{q}$$

— matches previous results of [McOrist-Melnikov](#); [Donagi-Guffin-Katz-ES](#)

— reduce to ordinary quantum cohomology on (2,2) locus

Example: Hirzebruch surfaces \mathbb{F}_n

2-pt correlation functions:

$$\langle \sigma \sigma \rangle = \alpha^{-1} [\Delta - \beta_1 \beta_2 \det(A + B) + (\gamma_1 + \beta_1)(\gamma_2 + \beta_2) \det B]$$

$$\langle \sigma \tilde{\sigma} \rangle = \alpha^{-1} \Delta$$

$$\langle \tilde{\sigma} \tilde{\sigma} \rangle = \alpha^{-1} [\Delta + \gamma_1 \gamma_2 \det(A + B) - (\gamma_1 + \beta_1)(\gamma_2 + \beta_2) \det A]$$

where

$$\Delta = \beta_1 \beta_2 \det A - \gamma_1 \gamma_2 \det B$$

$$\alpha = \Phi_1 \Phi_2$$

for

$$\Phi_i = \beta_i^2 \det A - \beta_i \gamma_i \gamma_{AB} + \gamma_i^2 \det B$$

$$\gamma_{AB} = \det(A + B) - \det A - \det B$$

JKG residue results match Cech cohomology computation.

For a general toric variety + deformation of tangent bundle,

$$\langle f(\vec{\sigma}) \rangle = \sum_{\mathfrak{m}_1, \dots \in \mathbb{Z}} \text{JKG} - \text{Res}_{\vec{\sigma}=0} \left\{ \left[\prod_{a, \alpha} \left(\frac{1}{\det M_{(\alpha)}} \right)^{Q_{\alpha}^a (\mathfrak{m}_a) + 1} q_a^{\mathfrak{m}_a} \right] f(\vec{\sigma}) \right\}$$

It can be shown that this matches result of [McOrist-Melnikov '08](#):

$$\langle f(\sigma) \rangle = \sum_{\sigma | J=0} f(\sigma) \left((\det J_{a,b}) \prod_{\alpha} \det M_{(\alpha)} \right)^{-1}$$

where
$$J_a = \ln \left(q_a^{-1} \prod_{\alpha} M_{(\alpha)}^{Q_{\alpha}^a} \right)$$

Now, getting new expressions for old results is nice, but, what's even better is that we can also get new results.....

Nonabelian cases

So far, we have discussed the results of applying susy localization to $A/2$ theories describing toric varieties.

Next: Grassmannians

Understanding $A/2$ twists of Grassmannians has been an open problem for many years, as older GLSM techniques don't easily apply.

We'll see that susy localization allows us to quickly derive results not previously obtainable.

Basic example, (2,2): $G(k,n)$ = Grassmannian of k planes in \mathbb{C}^n

Physics: $U(k)$ gauge theory

n chiral multiplets in fundamental rep'

(0,2) deformation:

$U(k)$ gauge theory

n chiral multiplets ϕ^i in fundamental rep'

n Fermi multiplets Λ^i in fundamental rep'

$$\bar{D}_+ \Lambda_a^i = \sigma_a^b \phi_b^i + B_j^i (\text{Tr } \sigma) \phi_a^j$$

The B 's define deformation off (2,2) locus.

Can show, total num' of deformations = $n^2 - 1$

(for $1 < k < n - 1$)

— overall trace of B defines trivial deformation; rest interesting

General formula for $A/2$ correlation functions:

$$\langle f(\sigma_1, \dots, \sigma_k) \rangle = \frac{1}{k!} \sum_{m_1, \dots, m_k \in \mathbb{Z}} \text{JKG} - \text{Res}_{\vec{\sigma}=0} \left\{ q^{\sum m_i} \left(\prod_{\alpha \neq \beta} (\sigma_\alpha - \sigma_\beta) \right) \prod_{\alpha=1}^k \left(\frac{1}{\det \tilde{E}(\sigma_\alpha)} \right)^{m_\alpha+1} f(\vec{\sigma}) \right\}$$

$$\text{where } \tilde{E}_j^i(\sigma) = \sigma \delta_j^i + B_j^i \left(\sum_{\alpha} \sigma_\alpha \right)$$

Q.s.c. relations: $\det \tilde{E}(\sigma_\alpha) = q$ for all α

We'll see more meaningful expressions shortly....

We'll give formulas for cases in which B is diagonal, for simplicity.

Example: Deformation of TG(2,4)

Classical correlation functions ($\mathbf{m}_1 = \mathbf{m}_2 = 0$)

$$\langle \sigma_1^4 \rangle = \Delta^{-1} (I_1 + 2I_1^2 + 4I_1I_2 - 2I_3 + 2I_2^2 + 2I_1I_3 - 4 \det B + 2I_2I_3 - 2I_1 \det B)$$

$$\langle \sigma_1^3 \sigma_2 \rangle = \Delta^{-1} (-1 - 3I_1 - 2I_1^2 - 3I_2 - 4I_1I_2 - 2I_2^2 - I_3 - 2I_1I_3 + 4 \det B - 2I_2I_3 + 2I_1 \det B)$$

$$\langle \sigma_1^2 \sigma_2^2 \rangle = \Delta^{-1} (2 + 4I_1 + 2I_1^2 + 4I_2 + 4I_1I_2 + 2I_3 - 4 \det B + 2I_2^2 + 2I_1I_3 + 2I_2I_3 - 2I_1 \det B)$$

$$\langle \sigma_1 \sigma_2^3 \rangle = \langle \sigma_1^3 \sigma_2 \rangle \quad \langle \sigma_2^4 \rangle = \langle \sigma_1^4 \rangle$$

$\Delta = 2 \prod_{i < j} (1 + B_{ii} + B_{jj})$ is the locus on which bundle degenerates.

$$I_1 = \sum_i B_{ii} = \text{tr } B \quad I_2 = \sum_{i < j} B_{ii} B_{jj} \quad I_3 = \sum_{i < j} B_{ii} B_{jj} B_{kk}$$

are coefficients in the characteristic polynomial of B .

How can we interpret those correlation functions usefully?
How can we compare to ordinary cohomology, on (2,2) locus?

Gauge-invariant combinations naturally correspond to Young diagrams (via 'Schur polynomials'):

$$\sigma_{\square} = \sigma_1 + \sigma_2$$


$$\sigma_{\square\square} = \sigma_1^2 + \sigma_2^2 + \sigma_1\sigma_2$$

$$\sigma_{\begin{array}{c} \square \\ \square \end{array}} = \sigma_1\sigma_2$$

$$\sigma_{\begin{array}{cc} \square & \square \\ \square & \end{array}} = \sigma_1^2\sigma_2 + \sigma_1\sigma_2^2$$

$$\sigma_{\begin{array}{cc} \square & \square \\ \square & \square \end{array}} = \sigma_1^2\sigma_2^2$$

Cohomology of $G(k,n)$ is naturally in 1-1 correspondence with Young diagrams inside $k \times (n-k)$ box.



so on $(2,2)$ locus, for example, $\sigma_{\square\square\square} = 0$
 $\sigma_{\begin{smallmatrix} \square & \square & \square \\ \square \end{smallmatrix}} = 0$

Classical correlation functions on $(2,2)$ locus:

$$\langle \sigma_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} \rangle = +1 \quad \langle \sigma_{\square\square\square\square} \rangle = 0 = \langle \sigma_{\begin{smallmatrix} \square & \square & \square \\ \square \end{smallmatrix}} \rangle$$

which imply OPE $\sigma_{\square\square\square} \cdot \sigma_{\square} = 0$

hence $\sigma_{\square\square\square} = 0$ Agree!

$$\begin{aligned}
(2,2) \text{ locus: } \quad \sigma_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} &= \sigma_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}} = \sigma_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} = 0 & \quad \text{classically} \\
\sigma_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}} - \sigma_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} &= 2q & \quad \text{nonpert'ly}
\end{aligned}$$

(0,2):

$$(a) \quad (1 + I_1 + I_2 + I_3)\sigma_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} + (I_3 + 2I_2 + 2I_1)\sigma_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array}} = 0$$

$$(b) \quad (1 + I_1 + I_2 + I_3)\sigma_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}} + (1 + 3I_1 + 3I_2 + 2I_3)\sigma_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} + (I_3 + 2I_2 + 2I_1)\sigma_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = 0$$

$$\begin{aligned}
(c) \quad (1 + I_3 + I_2 + I_1 + 2 \det B)\sigma_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}} &+ (-1 + I_2 + 3I_1 + 6 \det B)\sigma_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} \\
&+ (-I_3 + 2I_1 + 4 \det B)\sigma_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = 2q
\end{aligned}$$

(derived from $\det E = q$)

More generally, for any deformation of given form of $TG(k,n)$, we've recently argued that

classical sheaf cohomology ring =

$$\mathbb{C}[\sigma_{(1)}, \dots, \sigma_{(k(n-k))}] / (D_{k+1}, \dots, D_{k(n-k)}, R_{n-k+1}, \dots, R_{k(n-k)})$$

$$D_m = \det(\sigma_{1+j-i})_{1 \leq i, j \leq m}$$

$$R_r = \sum_{i=0}^{\min(r,n)} I_{n-i} \sigma_{(r-i)} \sigma_{(1)}^i$$

for $\sigma_{(1)} = \sigma_{\square}$ and so forth, and

(I_i) the coefficients in the characteristic polynomial of B .

$$\text{Exs: } I_0 = 1, \quad I_1 = \text{Tr } B, \quad I_n = \det B$$

More generally, for any deformation of given form of $TG(k,n)$, we've argued that

classical sheaf cohomology ring =

$$\mathbb{C}[\sigma_{(1)}, \dots, \sigma_{(k(n-k))}] / (D_{k+1}, \dots, D_{k(n-k)}, R_{n-k+1}, \dots, R_{k(n-k)})$$

$$\text{On } (2,2) \text{ locus, } R_r = \sigma_{(r)}$$

and the above becomes a standard presentation of the ordinary cohomology of $G(k,n)$.

So: matches $(2,2)$ locus.

Structure of quantum sheaf cohomology ring for a generic deformation of $T G(k,n)$

$$\mathbb{C}[\sigma_{(1)}, \sigma_{(2)}, \dots] / \langle D_{k+1}, D_{k+2}, \dots, R_{(n-k+1)}, \dots, R_{(n-1)}, \\ R_{(n)} + q, R_{(n+1)} + q\sigma_{(1)}, R_{(n+2)} + q\sigma_{(2)}, \dots \rangle$$

where $D_m = \det(\sigma_{(1+j-i)})_{1 \leq i, j \leq m}$

$$R_{(r)} = \sum_{i=0}^{\min(r,n)} I_i \sigma_{(r-i)} \sigma_{(1)}^i$$

for I_i the char' poly's of B: $\det(tI + B) = \sum_{i=0}^n I_{n-i} t^i$

Exs: $I_0 = 1, I_1 = \text{Tr } B, I_n = \det B$

Quantum sheaf cohomology ring:

$$\mathbb{C}[\sigma_{(1)}, \sigma_{(2)}, \dots] / \langle D_{k+1}, D_{k+2}, \dots, R_{(n-k+1)}, \dots, R_{(n-1)}, \\ R_{(n)} + q, R_{(n+1)} + q\sigma_{(1)}, R_{(n+2)} + q\sigma_{(2)}, \dots \rangle$$

If we turn off the deformation (set $B=0$), then

$$R_{(n)} = \sigma_{(n)}$$

and with some work it can be shown that the ring above can be presented as

$$\mathbb{C}[\sigma_{(1)}, \dots, \sigma_{(n-k)}] / \langle D_{k+1}, \dots, D_{n-1}, D_n + (-)^n q \rangle$$

which is a standard presentation of the (ordinary) quantum cohomology ring of $G(k,n)$.

(Buch, Kresch, Tamvakis, Bertram, Witten, Siebert, Tian,)

Example: $G(1,3)$

This has no nontrivial deformations, so any result should be equivalent to ordinary quantum cohomology ring of \mathbb{P}^2 .

$$\mathbb{C}[\sigma_{(1)}, \sigma_{(2)}, \dots] / \langle D_2, \dots, R_{(3)} + q, R_{(4)} + q\sigma_{(1)}, \dots \rangle$$

$$\text{which} = \mathbb{C}[\sigma_{(1)}] / \langle R_{(3)} + q \rangle$$

using $D_2 = \sigma_{(1)}^2 - \sigma_{(2)}, \dots$ to eliminate $\sigma_{(m)}$ for $m > 1$, and

$$\text{the result } R_{(3+\ell)} + q\sigma_{(\ell)} = \sigma_{(\ell)}(R_{(3)} + q)$$

$$\text{Now, } R_{(3)} = \sum_{i=0}^3 I_i \sigma_{(3-i)} \sigma^i = \left(\sum_{i=0}^3 I_i \right) \sigma^3 = (\det(I + B)) \sigma^3$$

so the qsc ring is $\mathbb{C}[\sigma] / \langle \det(I + B)\sigma^3 + q \rangle$

which is equivalent to std quantum cohomology ring.

So far we've used susy localization & other methods to

- find new residue-based formulas for toric results

- derive new results for nonabelian GLSM's
(Grassmannian quantum sheaf cohomology)

Next: Toda duals, then B/2 results

Toda duals

The mirror to the A model on \mathbb{P}^n is a B-twisted Landau-Ginzburg model, defined by a superpotential

$$W = X_1 + \cdots + X_n + \frac{q}{X_1 \cdots X_n}$$

often referred to as the 'Toda dual.'

Analogous statements are known for heterotic theories, which we'll describe, but first let's review how this works.

Toda duals (to \mathbb{P}^n)

$$W = X_1 + \cdots + X_n + \frac{q}{X_1 \cdots X_n}$$

Genus zero correlation functions:

$$\langle f(X_1, \cdots, X_n) \rangle = \sum_{d_{\ln X} W=0} \frac{f(X_1, \cdots, X_n)}{\det(\partial_{\ln X}^2 W)}$$

$$dW = 0 \implies X_1 = X_2 = \cdots = X_n \equiv X$$
$$\& X = qX^{-n} \text{ or } X^{n+1} = q \quad (\text{q.c. rel'n!})$$

Can show $\det(\partial^2 W) = (n+1)X^n$

hence $\langle X^m \rangle = \sum_{X^{n+1}=q} \frac{X^m}{(n+1)X^n}$

thus $\langle X^{n+d(n+1)} \rangle = q^d$ matching A model.

What's the heterotic analogue?

A heterotic Landau-Ginzburg model is defined by

- complex Kahler manifold X
- holomorphic vector bundle $\mathcal{E} \rightarrow X$
- holomorphic section $(J_a) \in \Gamma(\mathcal{E}^*)$

Recover ordinary Landau-Ginzburg models when

$$\mathcal{E} = TX, \quad J_a = \partial_a W$$

The mirror to the (A/2) theory on $\mathbb{P}^n \times \mathbb{P}^m$,
with def' of tangent bundle param'd by matrices A, B, C, D ,
is a Landau-Ginzburg theory on $(\mathbb{C}^\times)^n \times (\mathbb{C}^\times)^m$ with $\mathcal{E} = T$

$$J_i = a^{(1-n)/n} \left(aX_i + b \frac{\tilde{X}_1^{n+1}}{X_1^n} + \sum_{i=1}^n \mu_{n+1-i} \frac{\tilde{X}_1^i}{X_1^{i-1}} - \frac{q_1}{X_1 \cdots X_n} \right)$$

$$\tilde{J}_k = d^{(1-m)/m} \left(d\tilde{X}_k + c \frac{X_1^{m+1}}{\tilde{X}_1^m} + \sum_{k=1}^m \nu_k \frac{X_1^k}{\tilde{X}_1^{k-1}} - \frac{q_2}{\tilde{X}_1 \cdots \tilde{X}_m} \right)$$

$$a = \det A, \quad b = \det B, \quad c = \det C, \quad d = \det D,$$

$$\det(Ax + By) = ax^{n+1} + by^{n+1} + \sum_{i=1}^n \mu_i x^i y^{n+1-i},$$

$$\det(Cx + Dy) = cx^{m+1} + dy^{m+1} + \sum_{k=1}^m \nu_k x^k y^{m+1-k},$$

Let me outline correlation functions in these theories.

For heterotic LG models of the form just discussed,
at genus 0,

$$\langle f(X_i, \tilde{X}_k) \rangle = \sum_{J, \tilde{J}=0} \frac{f(X_i, \tilde{X}_k)}{\det(\partial(J, \tilde{J}))}$$

$$J, \tilde{J} = 0 \implies X_1 = \dots = X_n \equiv X, \quad \tilde{X}_1 = \dots = \tilde{X}_m \equiv \tilde{X}$$

$$\& \quad \det(AX + B\tilde{X}) = q, \quad \det(CX + D\tilde{X}) = \tilde{q}$$

— the quantum sheaf cohomology ring rel'ns

Can show all (genus 0) correlation functions match those of the corresponding $A/2$ theory, which is how we've checked this proposal.

Analogues for the B/2 model

So far I've only discussed susy localization in the A/2 model, for deformations of (2,2) theories.

We can also apply the same ideas to B/2 twists of 'dual' theories.

Which theories?

Recall mentioned earlier that

$$A/2(X, \mathcal{E}) = B/2(X, \mathcal{E}^*)$$

so we're going to be able to apply B/2 to spaces with deformations of cotangent bundles — no (2,2) locus.

Analogues for the B/2 model

Quick aside: how is this related to (0,2) mirror symmetry?

Suppose (0,2) NLSM's on (X, \mathcal{E}) and (Y, \mathcal{F}) define same SCFT.

$$\begin{array}{ccc} A/2(X, \mathcal{E}) & \xlongequal{\text{(0,2) mirror}} & B/2(Y, \mathcal{F}) \\ \parallel & & \parallel \\ B/2(X, \mathcal{E}^*) & \xlongequal{\text{(0,2) mirror}} & A/2(Y, \mathcal{F}^*) \end{array}$$

Analogues for the B/2 model

So, for example, we should be able to compute B/2 correlation functions for deformations of cotangent bundle of $\mathbb{P}^1 \times \mathbb{P}^1$.

Math:

$$0 \longrightarrow \mathcal{E} \longrightarrow \mathcal{O}(-1, 0)^2 \oplus \mathcal{O}(0, -1)^2 \xrightarrow{*} \mathcal{O}^2 \longrightarrow 0$$

$$* = \begin{bmatrix} Ax & Bx \\ C\tilde{x} & D\tilde{x} \end{bmatrix}$$

x, \tilde{x} vectors of homogeneous coordinates,
 A, B, C, D 2×2 matrices describing deformation

No (2,2) locus; but cotangent bundle at

$$A = D = I_{2 \times 2}, \quad B = C = 0$$

Physics....

Analogues for the B/2 model

So, for example, we should be able to compute B/2 correlation functions for deformations of cotangent bundle of $\mathbb{P}^1 \times \mathbb{P}^1$.

Physics:

x^i, \tilde{x}^i chiral superfields charge (1,0), (0,1)

p, \tilde{p} neutral chiral superfields

$\Lambda_i, \tilde{\Lambda}_i$ Fermi superfields charge (-1,0), (0,-1)

plus (0,2) superpotential

$$W = \Lambda_i F_j^i x^j + \tilde{\Lambda}_i \tilde{F}_j^i \tilde{x}^j$$

where $F_j^i = A_j^i p + B_j^i \tilde{p}$

$$\tilde{F}_j^i = C_j^i p + D_j^i \tilde{p}$$

(Compare A/2 version: there, no superpotential, and charges matched.)

Analogues for the B/2 model

So, for example, we should be able to compute B/2 correlation functions for deformations of cotangent bundle of $\mathbb{P}^1 \times \mathbb{P}^1$.

Unlike the A/2 case, here there is no σ field — no adjoint-valued scalar that is part of vector multiplet on (2,2) locus.

So, no Coulomb branch along which to compute.

Instead, have p field, which plays a `dual' role.

In effect, the Coulomb branch replaced by (part of) Higgs branch.

Analogues for the B/2 model

So, for example, we should be able to compute B/2 correlation functions for deformations of cotangent bundle of $\mathbb{P}^1 \times \mathbb{P}^1$.

Result:

Correlation functions are given by:

$$\langle f(p, \tilde{p}) \rangle = \sum_{m_1, m_2 \in \mathbb{Z}} \text{JKG} - \text{Res}_{p=\tilde{p}=0} \left\{ \left(\frac{1}{\det F} \right)^{m_1+1} \left(\frac{1}{\det \tilde{F}} \right)^{m_2+1} q^{m_1} \tilde{q}^{m_2} f(p, \tilde{p}) \right\}$$

$$\text{where } F = Ap + B\tilde{p} \quad \tilde{F} = Cp + D\tilde{p}$$

— equivalent to results in dual A/2 model, as expected

Other cotangent bundle deformations similar.

Summary

- Open problems in heterotic compactifications
- $A/2$, $B/2$ pseudo-topological field theories
- Susy localization in $A/2$ model for def's of $(2,2)$ theories
- Examples: $\mathbb{P}^1 \times \mathbb{P}^1$, \mathbb{F}_n , $G(k, n)$
 - new expressions for old results: JKG residues
 - new results: nonabelian GLSM's
- $(0,2)$ Toda duals
- Analogous computations in dual $B/2$ theories