#### CMB B-mode Polarization Experiments: First Results from POLARBEAR (and BICEP2)

#### Haruki Nishino (Kavli IPMU)

@ICRR/IPMU ACP seminar 2014-06-05

# "Major discovery!" by BICEP2

#### BICEP2 I: DETECTION OF B-mode POLARIZATION AT DEGREE ANGULAR SCALES

BICEP2 COLLABORATION - P. A. R. ADE<sup>1</sup>, R. W. AIKIN<sup>2</sup>, D. BARKATS<sup>3</sup>, S. J. BENTON<sup>4</sup>, C. A. BISCHOFF<sup>5</sup>, J. J. BOCK<sup>2,6</sup>, J. A. BREVIK<sup>2</sup>, I. BUDER<sup>5</sup>, E. BULLOCK<sup>7</sup>, C. D. DOWELL<sup>6</sup>, L. DUBAND<sup>8</sup>, J. P. FILIPPINI<sup>2</sup>, S. FLIESCHER<sup>9</sup>, S. R. GOLWALA<sup>2</sup>, M. HALPERN<sup>10</sup>, M. HASSELFIELD<sup>10</sup>, S. R. HILDEBRANDT<sup>2,6</sup>, G. C. HILTON<sup>11</sup>, V. V. HRISTOV<sup>2</sup>, K. D. IRWIN<sup>12,13,11</sup>, K. S. KARKARE<sup>5</sup>, J. P. KAUFMAN<sup>14</sup>, B. G. KEATING<sup>14</sup>, S. A. KERNASOVSKIY<sup>12</sup>, J. M. KOVAC<sup>5,17</sup>, C. L. KUO<sup>12,13</sup>, E. M. LEITCH<sup>15</sup>, M. LUEKER<sup>2</sup>, P. MASON<sup>2</sup>, C. B. NETTERFIELD<sup>4,16</sup>, H. T. NGUYEN<sup>6</sup>, R. O'BRIENT<sup>6</sup>, R. W. OGBURN IV<sup>12,13</sup>, A. ORLANDO<sup>14</sup>, C. PRYKE<sup>9,7,17</sup>, C. D. REINTSEMA<sup>11</sup>, S. RICHTEF<sup>5</sup>, R. SCHWARZ<sup>9</sup>, C. D. SHEEHY<sup>9,15</sup>, Z. K. STANISZEWSKI<sup>2,6</sup>, R. V. SUDIWALA<sup>1</sup>, G. P. TEPLY<sup>2</sup>, J. E. TOLAN<sup>12</sup>, A. D. TURNER<sup>6</sup>, A. G. VIEREGG<sup>5,15</sup>, C. L. WONG<sup>5</sup>, AND K. W. YOON<sup>12,13</sup>

to be submitted to a journal TBD

#### ABSTRACT

We report results from the BICEP2 experiment, a Cosmic Microwave Background (CMB) polarimeter specifically designed to search for the signal of inflationary gravitational waves in the B-mode power spectrum around  $\ell \sim 80$ . The telescope comprised a 26 cm aperture all-cold refracting optical system equipped with a focal plane of 512 antenna coupled transition edge sensor (TES) 150 GHz bolometers each with temperature sensitivity of  $\approx 300 \ \mu K_{CMB} \sqrt{s}$ . BICEP2 observed from the South Pole for three seasons from 2010 to 2012. A low-foreground region of sky with an effective area of 380 square degrees was observed to a depth of 87 nK-degrees in Stokes Q and U. In this paper we describe the observations, data reduction, maps, simulations and results. We find an excess of B-mode power over the base lensed-ACDM expectation in the range  $30 < \ell < 150$ , inconsistent with the null hypothesis at a significance of >  $5\sigma$ . Through jackknife tests and simulations based on detailed calibration measurements we show that systematic contamination is much smaller than the observed excess. We also estimate potential foreground signals and find that available models predict these to be considerably smaller than the observed signal. These foreground models possess no significant cross-correlation with our maps. Additionally, cross-correlating BICEP2 against 100 GHz maps from the BICEP1 experiment, the excess signal is confirmed with  $3\sigma$  significance and its spectral index is found to be consistent with that of the CMB, disfavoring synchrotron or dust at 2.3 $\sigma$  and 2.2 $\sigma$ , respectively. The observed B-mode power spectrum is wellfit by a lensed-ACDM + tensor theoretical model with tensor/scalar ratio  $r = 0.20^{+0.07}_{-0.05}$ , with r = 0 disfavored at 7.0 $\sigma$ . Subtracting the best available estimate for foreground dust modifies the likelihood slightly so that r = 0is disfavored at 5.9 $\sigma$ .

#### You might already know well the BICEP2 result... Any progress so far after then?

#### Just a week before the BICEP2 paper...

#### A MEASUREMENT OF THE COSMIC MICROWAVE BACKGROUND B-MODE POLARIZATION POWER SPECTRUM AT SUB-DEGREE SCALES WITH POLARBEAR

THE POLARBEAR COLLABORATION: P.A.R. ADE<sup>29</sup>, Y. AKIBA<sup>32</sup>, A.E. ANTHONY<sup>2,5</sup>, K. ARNOLD<sup>14</sup>, M. ATLAS<sup>14</sup>, D.
BARRON<sup>14</sup>, D. BOETTGER<sup>14</sup>, J. BORRILL<sup>3,31</sup>, S. CHAPMAN<sup>9</sup>, Y. CHINONE<sup>17,13</sup>, M. DOBBS<sup>25</sup>, T. ELLEFLOT<sup>14</sup>, J. ERRARD<sup>31,3</sup>,
G. FABBIAN<sup>1,18</sup>, C. FENG<sup>14</sup>, D. FLANIGAN<sup>13,10</sup>, A. GILBERT<sup>25</sup>, W. GRAINGER<sup>28</sup>, N.W. HALVERSON<sup>2,5,15</sup>, M. HASEGAWA<sup>17,32</sup>,
K. HATTORI<sup>17</sup>, M. HAZUMI<sup>17,32,20</sup>, W.L. HOLZAPFEL<sup>13</sup>, Y. HORI<sup>17</sup>, J. HOWARD<sup>13,16</sup>, P. HYLAND<sup>24</sup>, Y. INOUE<sup>32</sup>, G.C.
JAEHNIG<sup>2,15</sup>, A.H. JAFFE<sup>11</sup>, B. KEATING<sup>14</sup>, Z. KERMISH<sup>12</sup>, R. KESKITALO<sup>3</sup>, T. KISNER<sup>3,31</sup>, M. LE JEUNE<sup>1</sup>, A.T. LEE<sup>13,27</sup>,
E.M. LEITCH<sup>4,19</sup>, E. LINDER<sup>27</sup>, M. LUNGU<sup>13,8</sup>, F. MATSUDA<sup>14</sup>, T. MATSUMURA<sup>17</sup>, X. MENG<sup>13</sup>, N.J. MILLER<sup>22</sup>, H. MORII<sup>17</sup>,
S. MOYERMAN<sup>14</sup>, M.J. MYERS<sup>13</sup>, M. NAVAROLI<sup>14</sup>, H. NISHINO<sup>20</sup>, H. PAAR<sup>14</sup>, J. PELOTON<sup>1</sup>, D. POLETTI<sup>1</sup>, E. QUEALY<sup>13,26</sup>,
G. REBEIZ<sup>6</sup>, C.L. REICHARDT<sup>13</sup>, P.L. RICHARDS<sup>13</sup>, C. ROSS<sup>9</sup>, I. SCHANNING<sup>14</sup>, D.E. SCHENCK<sup>2,5</sup>, B.D. SHERWIN<sup>13,21</sup>, A. SHIMIZU<sup>32</sup>, C. SHIMMIN<sup>13,7</sup>, M. SHIMON<sup>30,14</sup>, P. SIRITANASAK<sup>14</sup>, G. SMECHER<sup>33</sup>, H. SPIELER<sup>27</sup>, N. STEBOR<sup>14</sup>, B.
STEINBACH<sup>13</sup>, R. STOMPOR<sup>1</sup>, A. SUZUKI<sup>13</sup>, S. TAKAKURA<sup>23,17</sup>, T. TOMARU<sup>17</sup>, B. WILSON<sup>14</sup>, A. YADAV<sup>14</sup>, O. ZAHN<sup>27</sup>

Draft version March 10, 2014

#### ABSTRACT

We report a measurement of the *B*-mode polarization power spectrum in the cosmic microwave background (CMB) using the POLARBEAR experiment in Chile. The faint *B*-mode polarization signature carries information about the Universe's entire history of gravitational structure formation, and the cosmic inflation that may have occurred in the very early Universe. Our measurement covers the angular multipole range  $500 < \ell < 2100$  and is based on observations of  $30 \deg^2$  with 3.5'resolution at 150 GHz. On these angular scales, gravitational lensing of the CMB by intervening structure in the Universe is expected to be the dominant source of *B*-mode polarization. Including both systematic and statistical uncertainties, the hypothesis of no *B*-mode polarization power from gravitational lensing is rejected at 97.5% confidence. The band powers are consistent with the standard cosmological model. Fitting a single lensing amplitude parameter  $A_{BB}$  to the measured band powers,  $A_{BB} = 1.12 \pm 0.61(\text{stat})^{+0.04}_{-0.10}(\text{sys}) \pm 0.07(\text{multi})$ , where  $A_{BB} = 1$  is the fiducial WMAP-9 ACDM value. In this expression, "stat" refers to the statistical uncertainty, "sys" to the systematic uncertainty associated with possible biases from the instrument and astrophysical foregrounds, and "multi" to the calibration uncertainties that have a multiplicative effect on the measured amplitude  $A_{BB}$ .

#### POLARBEAR published the best B-mode measurement at subdegree scale!

#### First result from ACTpol is out too.

#### THE ATACAMA COSMOLOGY TELESCOPE: CMB POLARIZATION AT $200 < \ell < 9000$

SIGURD NAESS<sup>1</sup>, MATTHEW HASSELFIELD<sup>2,3</sup>, JEFF MCMAHON<sup>4</sup>, MICHAEL D. NIEMACK<sup>5</sup>, GRAEME E. ADDISON<sup>3</sup>,
PETER A. R. ADE<sup>6</sup>, RUPERT ALLISON<sup>1</sup>, MANDANA AMIRI<sup>3</sup>, ANDREW BAKER<sup>7</sup>, NICK BATTAGLIA<sup>8</sup>, JAMES A. BEALL<sup>9</sup>,
FRANCESCO DE BERNARDIS<sup>5</sup>, J RICHARD BOND<sup>10</sup>, JOE BRITTON<sup>9</sup>, ERMINIA CALABRESE<sup>1</sup>, HSIAO-MEI CHO<sup>9</sup>,
KEVIN COUGHLIN<sup>4</sup>, DEVIN CRICHTON<sup>11</sup>, SUDEEP DAS<sup>12</sup>, RAHUL DATTA<sup>4</sup>, MARK J. DEVLIN<sup>13</sup>, SIMON R. DICKER<sup>13</sup>,
JOANNA DUNKLEY<sup>1</sup>, ROLANDO DÜNNER<sup>14</sup>, JOSEPH W. FOWLER<sup>9</sup>, ANNA E. FOX<sup>9</sup>, PATRICIO GALLARDO<sup>5,14</sup>, EMILY GRACE<sup>15</sup>,
MEGAN GRALLA<sup>11</sup>, AMIR HAJIAN<sup>10</sup>, MARK HALPERN<sup>3</sup>, SHAWN HENDERSON<sup>5</sup>, J. COLIN HILL<sup>2</sup>, GENE C. HILTON<sup>9</sup>,
MATT HILTON<sup>16</sup>, ADAM D. HINCKS<sup>3</sup>, RENÉE HLOZEK<sup>2</sup>, PATTY HO<sup>15</sup>, JOHANNES HUBMAYR<sup>9</sup>, KEVIN M. HUFFENBERGER<sup>17</sup>,
JOHN P. HUGHES<sup>7</sup>, LEOPOLDO INFANTE<sup>14</sup>, KENT IRWIN<sup>18</sup>, REBECCA JACKSON<sup>4,19</sup>, JEFF KLEIN<sup>13</sup>, BRIAN KOOPMAN<sup>5</sup>,
ARTHUR KOSOWSKY<sup>20</sup>, DALE LI<sup>9</sup>, THIBAUT LOUIS<sup>1</sup>, MARIUS LUNGU<sup>13</sup>, MATHEW MADHAVACHERIL<sup>21</sup>,
TOBIAS A. MARRIAGE<sup>11</sup>, LOIC MAURIN<sup>14</sup>, FELIPE MENANTEAU<sup>22,23</sup>, KAVILAN MOODLEY<sup>16</sup>, CHARLES MUNSON<sup>4</sup>,
LAURA NEWBURGH<sup>15</sup>, JOHN NIBARGER<sup>9</sup>, MICHAEL R. NOLTA<sup>10</sup>, LYMAN A. PAGE<sup>15</sup>, CHRISTINE PAPPAS<sup>15</sup>,
BRUCE PARTRIDGE<sup>24</sup>, FELIPE ROJAS<sup>14</sup>, BENJAMIN SCHMITT<sup>13</sup>, NEELIMA SEHGAL<sup>21</sup>, BLAKE D. SHERWIN<sup>25</sup>, JON SIEVERS<sup>26,10</sup>,
SARA SIMON<sup>15</sup>, DAVID N. SPERGEL<sup>2</sup>, SUZANNE T. STAGGS<sup>15</sup>, ERIC R. SWITZER<sup>27,19</sup>, ROBERT THORNTON<sup>28,13</sup>, HY TRAC<sup>8</sup>, CAROLE TUCKER<sup>6</sup>, ALEXANDER VAN ENGEL<sup>21</sup>, JON WARD<sup>13</sup>, EDWARD J. WOLLACK<sup>27</sup>

Draft: May 22, 2014

#### ABSTRACT

We report on measurements of the cosmic microwave background (CMB) and celestial polarization at 146 GHz made with the Atacama Cosmology Telescope Polarimeter (ACTPol) in its first three months of observing. Four regions of sky covering a total of 270 square degrees were mapped with an angular resolution of 1.3'. The map noise levels in the four regions are between 11 and 17  $\mu$ K-arcmin. We present TT, TE, EE, TB, EB, and BB power spectra from three of these regions. The observed E-mode polarization power spectrum, displaying six acoustic peaks in the range 200 <  $\ell$  < 3000, is an excellent fit to the prediction of the best-fit cosmological models from WMAP9+ACT and *Planck* data. The polarization power spectrum, which mainly reflects primordial plasma velocity perturbations, provides an independent determination of cosmological parameters consistent with those based on the temperature power spectrum, which results mostly from primordial density perturbations. We find that without masking any point sources in the EE data at  $\ell$  < 9000, the Poisson tail of the EE power spectrum due to polarized point sources has an amplitude less than 2.4  $\mu$ K<sup>2</sup> at  $\ell$  = 3000 at 95% confidence. Finally, we report that the Crab Nebula, an important polarization calibration source at microwave frequencies, has 8.7% polarization with an angle of 150.9° ± 0.5° when smoothed with a 5'

- Only three months of data.
- High-ell E-mode power spectrum measurement
  - B-mode power spectrum is null consistent.

### First Results from POLARBEAR

- "A Measurement of the Cosmic Microwave Background B-Mode Polarization Power Spectrum at Sub-Degree Scales with POLARBEAR" (<u>http://arxiv.org/abs/1403.2369</u>)
  - Best measurement of B-mode auto-power spectrum in subdegree angular scale
- "Gravitational Lensing of Cosmic Microwave Background Polarization" (<u>http://arxiv.org/abs/1312.6646</u>, accepted at PRL)
  - First evidence for gravitational lensing with CMB polarization data alone
- "Evidence for Gravitational Lensing of the Cosmic Microwave Background Polarization from Cross-correlation with the Cosmic Infrared Background" (PRL 112, 131302 (2014))
  - Cross-correlation with CIB, confirmation of SPTpol result

Measurement of lensing B-modes with three independent methods

# **Cosmology with CMB B-modes**

- Primordial Gravitational Wave from Inflation
  - GUT-scale ultra high energy physics
- Weak Gravitational Lensing from Large Scale Structure
  Sum of neutrino mass

### **Cosmic Microwave Background**



Physics in the early universe is imprinted in CMB.

# Thomson scattering makes linear polarization



Wayne Hu (<u>http://background.uchicago.edu/</u> ~whu/intermediate/Polarization/polar1.html)

#### Local quadrupole anisotropy+Thomson scattering induced CMB polarization



Wayne Hu (<u>http://background.uchicago.edu/</u> ~whu/intermediate/Polarization/polar1.html)

### Density-wave (scalar perturbation)

A density wave produces temperature anisotropy.



### Density-wave (scalar perturbation)

Polarization pattern produced by the density wave





- Polarization fields can be decomposed into two orthogonal fields of E- and B-modes
  - Density perturbations (which have been seen as temperature fluctuation) can only create E-mode.

### Inflation



- Accelerating expansion (~e<sup>60</sup>) in 10<sup>-35</sup> seconds
- Two types of perturbations were generated during inflation from quantum fluctuations
  - Scalar (density) perturbations
  - Tensor perturbations = Gravitational wave

#### Gravitational wave (tensor perturbations)



#### Local quadrupole from gravitational wave

Gravitational wave produces local quadrupole temperature anisotropies

wave vector k



#### Polarization pattern by gravitational wave

Polarization pattern generated by the quadrupole

wave vector k



#### Polarization pattern by gravitational wave



#### Polarization pattern by gravitational wave

Let's consider the polarization projection on to the last scattering surface like this.



### E-mode from gravitational wave

wave vector k (projected)



"x" mode

### B-mode from gravitational wave

wave vector k (projected)



**B-mode:** Polarization

orientation is 45 or 135

rotated to the wave vector, k.





# **Scalar/Tensor Perturbations**

- Scalar perturbations:
  - Produces CMB Temperature fluctuation (and Emode polarization)
- Tensor perturbations: Gravitational Wave
  - Tiny effect in T or E
  - But produces B-mode polarization too
  - Unique prediction from inflation!

#### CMB power spectra



#### Quantitative Study of Inflation w/ B-mode

- Energy scale of inflation directly relates to the ratio of tensor and scalar perturbations  $A = (A + A)^{n_s - 1} + (A + A)^{n_t}$ 
  - Tensor-to-scalar ratio:

$$r \equiv \frac{A_t}{A_s} \qquad P_s = A_s \left(\frac{k}{k_0}\right)^{n_s - 1} \quad P_t = A_t \left(\frac{k}{k_0}\right)^{n_t}$$

- Proportional to B-mode power spectrum
- Inflation potential:  $V^{1/4} \sim (r/0.01)^{1/4} \times 10^{16} \text{ GeV}$
- Detecting B-mode in large angular scales proves inflation and enables quantitative study of models of inflation.
  - Also a probe of physics of ultra high energy scale (~GUT scale)
  - r=0.2  $\rightarrow$  energy scale of inflation: 2 x 10<sup>16</sup> GeV

#### Another B-mode source: Gravitational Weak Lensing





Observed CMB has been deflected by gravitational potentials of Large Scale Structure from z~1100 up to now.

### **CMB** Polarization and Weak Lensing



- Fig. from astro-ph/0111606
   Gravitational lensing mixes E-mode and B-mode.
- Primary B-mode source in small angular scale (subdegree).

### **CMB** Polarization and Weak Lensing

**Observed CMB polarization**  

$$(Q + iU)(\hat{\mathbf{n}}) = (\tilde{Q} + i\tilde{U})(\hat{\mathbf{n}} + \mathbf{d}(\hat{\mathbf{n}}))$$
  
deflection angle  
~: primordial CMB  
(weak lensing effect)

- Typical deflection angle: a few arcmin
- Coherent over ~2 deg. scale

Requires degree-scale observation with a telescope of arc minute resolution

### Weak Lensing and Neutrino Mass

Massive neutrinos suppress the evolution of large scale structures.



Higher neutrino mass ↓ Less large-scale structure ↓ Smaller lensing effect

Planck 2013 (lensing) w/ WMAP pol.+SPT+ACT  $\Sigma(m_v) < 0.85 \text{ eV} (95\% \text{CL})$ 

# Upper limits of B-mode power spectrum before March 2014

From BICEP1-3yr paper arXiv:1310.1422

27



#### Large angular scale Small angular scale

 SPTpol claimed the "detection of B-mode" by the cross-correlation with galaxy, but BB auto-correlation spectrum from SPTpol is not published yet.

# CMB polarization experiments

# **CMB** polarization experiments

POLARBEAR

- Satellite: PLANCK
- Balloon-borne: EBEX, SPIDER
- South Pole
  - BICEP and Keck Array
  - SPTpol
- Atacama Desert, Chile
  - POLARBEAR
  - ACTpol
  - Atacama B-mode Search (ABS)





ABS

ACT

# POLARBEAR



- Ground-based CMB polarization experiment in the Atacama desert
  - 5,200m altitude
  - Thin and dry atmosphere

#### Started observation since early 2012



International collaboration from 5countries, ~70 researchers

### **POLARBEAR site**

#### Northern Chile

US Dept of State Geographer Data SIO, NOAA, U.S. Navy, NGA, GEBCO © 2014 Google © 2014 Inav/Geosistemas SRL

Google earth

### **POLARBEAR site**

POLARBEAR

自然又几日

・リオハ州

アルセンチン

サルタ州

サンティアゴ・デル・エスデロ州

コルドバ州

#### Atacama Desert

デジー首都州 サンディアゴ US Dept of State Geographer Data SIO, NOAA, U.S. Navy, NGA, GEBGO Image Landsat\_サー © 2014 Inav Geosistemas SRL

サンファンル

699 km

33

一手儿

リオグランテ

マットグロッソ

パラグアイ

チャコ州

サンタフェ州

シトレ・リオス州 ラルクアイ

アスシンオンセントラル

ゴリエンテス州

Sooglezearth

### Atacama desert

ASTE, NANTEN ~4860m

POLARBEAR

TAO ~5640m (world highest)

TAO

APEX

ALMA

ALMA (QUIET) ~5000m

US Dept of State Geographer © 2014 Maperty

Image © 2014 DigitalClobe

2663 m

POLARBEAR, ACT, ABS ~5200m

Google earth

# Huan Tran Telescope



- Off-axis Gregoriane Dragone
- 3.5m Primary mirror
  - 3.5 arcmin (FWHM) resolution
  - designed to measure both primordial and lensing Bmodes

Receiver cryostat

# **POLARBEAR Receiver**



Focal Plane (cooled to 0.25K)

HWP (polarization modulation)

# Focal Plane Detector Array



- 2 bolometers in a pixel
- 637 pixels in a array
- 1274 bolometers (~80% yield)

Noise level: 550 uKs<sup>1/2</sup> /bolo 23 uKs<sup>1/2</sup> /array



# Polarization measurement by pair-differencing



Pair bolometers in a single pixel are sensitive to linear polarizations, which are perpendicular to each other.

Pixel differencing gives us a polarization information of incident light. It is also effective for common mode noise rejection.

Good beam (and gain) matching between pair detectors is crucial because the mismatch can potentially cause leakage from T->P.

# **BICEP/Keck Array**



- CMB polarization experiments at South Pole
  - Target large angular scales, i.e. the primordial gravitational wave B-modes

### Spec. Summary of BICEP/Keck

	BICEP1	BICEP2	Keck Array ≒	BICEP3
Operation Time	2006-2008	2010-2012	2011-	2014
Frequency	100/150 GHz	150 GHz	150GHz (100GHz)	100GHz
Detectors	98 NTD Ge	512 TES bolometers	2560 TES bolometers	2560 TES bolometers
Noise level (uK s	54 (/array)	16 (/array)	~10 (/array)	
Beam (Resolution)	0.93°/0.60° FWHM	0.52° FWHM	0.52° FWHM	0.37° FWHM
comparable to SPTpol, POLARBEAR, etc			optimized for large angular scale	
BICEP/Keck also empolys similar detector technique (TES bolometers + pair-differencing).				

### **Observed Sky Region**

Planck 857GHz map



- Choose low astrophysical "foreground" regions
- POLARBEAR: 3 small sky regions (3°x3°): ~30 deg<sup>2</sup>
- **BICEP**: ~380 deg<sup>2</sup>, fsky~2%

### Telescope scan

- Always scanning in azimuth (horizontally) at each constant elevation
  - To see the modes before detector gain drifting.

# **CMB** Data Analysis

- Calibration: Gain, Pointing, Beam, Polarization angle...
- Data Selection: Bad weather, Badly tuned detectors...
- Map-making: I, Q, U maps of each field (for each day)
- Power spectrum: Cross-spectrum with each-day-maps
- Null Tests (Jackknife): data validation, systematics check
- Systematic Error: Estimate leakage from T,  $E \rightarrow B$

### **Beam Measurement**

#### POLARBEAR

#### **BICEP2**



### **CMB** Polarization Maps (POLARBEAR)



- (Filtered) map noise level: 5 uK-arcmin
- ~10 times deeper than Planck
  - Deepest map in this scale ~ BICEP2

### **BICEP2 CMB Maps**



RMS noise level of 87 nK in 1° x 1° pixel

### BICEP2 E-mode & B-mode maps



Impressive demonstration of E and B modes. We can see B-modes in map space.

## Null Tests (Jackknife)



Split the data into two with some criteria to illuminate some possible systematic effect

Subtract spectra with each other and Check if the residual spectrum is null or not.

- If there is a systematic bias between the two data set, the difference spectrum would not be null.
- BICEP2 did this in a map space.

# Null Tests (POLARBEAR)

- First half vs. Second half
- Rising vs. Setting
- High vs. Low elevation
- High vs. Low gain
- Good vs. Bad weather
- Pixel polarization angle orientation type
- Left vs Right side detectors
- Left vs. right-going scan
- Moon distance

### Null Test PTEs

#### POLARBEAR

#### **BICEP2** PTE (each spectrum)



No significant systematics found

### (After passing all the null tests...) B-mode power spectra

(BB power spectrum was blinded for POLARBEAR.)

#### Comparison with other upper limits (3/10)



#### Comparison with other upper limits (3/17)



# **BICEP2 BB spectrum**



- Very low PTE with the solid line (lensed ΛCDM)
  - 5.3σ detection of the excess power at low-ell
- What about the excess around ell~200?
  - "We caution against over interpretation of the two high band powers at ell~220 — their joint significance is less than 3σ"

These might be just statistical fluctuations, or might be an indication for unknown systematics?

### Cross-spectrum w/ Keck Array



If there are any systematics only in BICEP2 data, the cross correlation of BICEP2xKeck should show significantly different result.

- The cross-spectrum with Keck Array shows better fit with lensed ACDM+r>0 although it looks significantly different from the BICEP2 auto-spectrum.
  - This adds some confidence for non-zero r detection.
  - But this might also indicate unknown systematic contamination of BICEP2.

### **Constraint on r from BICEP2**



r=0.20<sup>+0.07</sup><sub>-0.05</sub>, 7.0σ r>0 detection

- w/ foreground subtraction (DDM2): r=0.16<sup>+0.06</sup>-0.05, 5.9σ
   detection
- (Best-fit r ranges between 0.12 ~ 0.21 depending on the model or the treatment of foreground)

## Then, what is going on now?

# Some worrying systematics

- Instrumental systematics
- Foreground contamination

#### Most worrying (instrumental) systematics for BICEP2: Differential Beam Systematics



 Pointing mismatch: Differential pointing can cause a serious leakage from temperature fluctuation to Bmode polarization.

Other differential beam effects





Differential beam systematics and "deprojection"

- Differential beam effect can be mitigated by "deprojection" technique in the BICEP2 data analysis using Planck T map.
- Without the deprojection, observed CIBB would be dominated by the leakage from CITT by differential pointing and relative gain mismatch.

Waiting for details in the coming systematics paper...

#### Multipole

10-7



### **Systematic** uncertainty (BICEP2)

**Every systematic** uncertainty is estimated to be negligible.

Waiting for details in the coming systematics paper...

#### Multipole

#### Systematic uncertainty (POLARBEAR)



Estimated foreground is much smaller compared with either stat or sys error.

pointing differential beam size polarization angle differential ellipticity relative gain (HWP dependent) relative gain electrical crosstalk gain drift

#### Negligible leakage without deprojection

### Astrophysical foregrounds (BICEP2)



- Dashed lines: autospectra of models
- Solid lines: crossspectra with BICEP2 maps

• DDM: "Data Driven Model" constructed from publicly available Planck data products. The Planck dust model map at 353GHz is scaled to 150 GHz.

#### What is wrong with this?

#### **Polarization Fraction**

Apparent polarization fraction (p) at 353 GHz, 1° resolution Not CIB subtracted  $\sqrt{O^2 + U^2}$ 

It seems BICEP2 used the Planck map image presented at a conference by digitizing it.

They estimated the polarization fraction to be 5%.

However, this map is not CIB (cosmic infrared background) subtracted, which could result in underestimating pol. fraction.

0.20

p ranges from 0 to ~20% Low p values in inner MW plane. Consistent with unpolarized CIB Large p values in outer plane and intermediate latitudes

0%

#### **Recent Planck dust polarization fraction map**



No one knows the answer... Wait for Planck's next release.

# Other analysis results from POLARBEAR

#### **Optimal Estimator for Gravitational Lensing**

- Gravitational lensing induces correlation between different multipoles (*I* and *I*') of E-mode and B-mode.
  - We can make use of the correlation to estimate the lensing deflection field, d(L).



arXiv:1312.6646  $C_{L}^{dd} = < dx dx'*>$ 

"4-point correlation" with only CMB polarization data alone PRL 113, 131302 (2014)  $C_L^{dI} = \langle dx, CIB \rangle$ 

Cross-correlation with CIB (biased tracer of dark matter halos)

## Combined lensing power spectra



- Consistent between <EEEB> and <EBEB>.
- 4.2 σ rejection of null hypothesis
- First detection (evidence) for gravitational lensing with CMB polarization alone

### **Cross-correlation with CIB**





CIB: Herschel/SPIRE ~500um

CIB is a good tracer for the matter distribution in  $z=1\sim3$ .  $\rightarrow$  Should correlate with CMB lensing.

- Reject null hypothesis at 4.0 σ
- Consitent with SPTpol (Hanson et al, 2013)

# POLARBEAR-2 & Simons Array

New POLARBEAR-2 receiver developed by KEK



Multichroic antenna(90+150GHz) +TES bolometer

#### Two new telescopes (Simons Array)



- POLARBEAR-2 will be deployed in 2015
- Confirm and measure r. Characterize B-mode better
- Constrain sum of neutrino mass:  $\sigma \sim 20$  meV (w/o sys.)

## Summary

- BICEP2 detected B-mode power at degree angular scale
  - Detection of non-zero r at > 5 σ, r=0.1~0.2
  - Recent discussions on foreground suggest that the detection of the cosmological signal is not conclusive.
    - Needs confirmations from other experiments!
- POLARBEAR measured B-mode power spectrum with the best sensitivity at sub-degree angular scale.
  - Detected gravitational lensing with three independent methods (at >  $4\sigma$ )
  - POLARBEAR is starting large angular scale observations.

#### "New era of B-mode cosmology has begun."